

Mutational Spectrum in a Worldwide Study of 29,700 Families with *BRCA1* or *BRCA2* Mutations

Timothy R. Rebbeck¹, Tara M. Friebe¹, Eitan Friedman², Ute Hamann³, Dezheng Huo⁴, Ava Kwong⁵, Edith Olah⁶, Olufunmilayo I. Olopade⁴, Angela R. Solano⁷, Soo-Hwang Teo⁸, Mads Thomassen⁹, Jeffrey N. Weitzel¹⁰, TL Chan MBBS¹¹, Fergus J. Couch¹², David E. Goldgar¹³, Torben A. Kruse⁹, Edenir Inêz Palmero¹⁴, Sue Kyung Park¹⁵, Diana Torres^{3,16}, Elizabeth J. van Rensburg¹⁷, Lesley McGuffog¹⁸, Michael T. Parsons¹⁹, Goska Leslie MEng¹⁸, Cora M. Aalfs²⁰, Julio Abugattas²¹, Julian Adlard²², Simona Agata²³, Kristiina Aittomäki²⁴, Lesley Andrews²⁵, Irene L. Andrulis²⁶, Adalgeir Arason²⁷, Norbert Arnold²⁸, Banu K. Arun²⁹, Ella Asseryanis³⁰, Leo Auerbach³⁰, Jacopo Azzollini³¹, Judith Balmaña³², Monica Barile³³, Rosa B. Barkardottir³⁴, Daniel Barrowdale¹⁸, Javier Benitez³⁵, Andreas Berger³⁶, Raanan Berger³⁷, Amie M. Blanco³⁸, Kathleen R. Blazer¹⁰, Marinus J. Blok³⁹, Valérie Bonadona⁴⁰, Bernardo Bonanni³³, Angela R. Bradbury⁴¹, Carole Brewer⁴², Bruno Buecher⁴³, Sandra S. Buys⁴⁴, Trinidad Caldes⁴⁵, Almuth Caliebe⁴⁶, Maria A. Caligo⁴⁷, Ian Campbell⁴⁸, Sandrine Caputo⁴³, Jocelyne Chiquette⁴⁹, Wendy K. Chung⁵⁰, Kathleen B.M. Claes⁵¹, J. Margriet Collée⁵², Jackie Cook⁵³, Rosemarie Davidson⁵⁴, Miguel de la Hoya⁴⁵, Kim De Leeneer⁵¹, Antoine de Pauw⁴³, Capucine Delnatte⁵⁵, Orland Diez⁵⁶, Yuan Chun Ding⁵⁷, Nina Ditsch⁵⁸, Susan M. Domchek⁴¹, Cecilia M. Dorfling MSc¹⁷, Carolina Velazquez⁵⁹, Bernd Dworniczak⁶⁰, Jacqueline Eason⁶¹, Douglas F. Easton¹⁸, Ros Eeles⁶², Hans Ehrencrona⁶³, Bent Ejlersen⁶⁴, EMBRACE¹⁸, Christoph Engel⁶⁵, Stefanie Engert⁶⁶, D. Gareth Evans⁶⁷, Laurence Faivre⁶⁸, Lidia Feliubadaló⁶⁹, Sandra Fert Ferrer⁷⁰, Lenka Foretova⁷¹, Jeffrey Fowler⁷², Debra Frost¹⁸, Henrique C. R. Galvão⁷³, Patricia A. Ganz⁷⁴, Judy Garber⁷⁵, Marion Gauthier-Villars⁴³, Andrea Gehrig⁷⁶, GEMO Study Collaborators⁷⁷, Anne-Marie Gerdes⁷⁸, Paul Gesta⁷⁹, Giuseppe Giannini⁸⁰, Sophie Giraud⁸¹, Gord Glendon⁸², Andrew K. Godwin⁸³, Mark H. Greene⁸⁴, Jacek Gronwald⁸⁵, Angelica Gutierrez-Barrera²⁹, Eric Hahnen⁸⁶, Jan Hauke⁸⁶, HEBON⁸⁷, Alex Henderson⁸⁸, Julia Hentschel⁸⁹, Frans B.L. Hogervorst⁹⁰, Ellen Honisch⁹¹, Evgeny N. Imyanitov⁹², Claudine Isaacs⁹³, Louise Izatt⁹⁴, Angel Izquierdo⁹⁵, Anna Jakubowska⁸⁵, Paul James⁹⁶, Ramunas Janavicius⁹⁷, Uffe Birk Jensen⁹⁸, Esther M. John⁹⁹, Vijai Joseph¹⁰⁰, Katarzyna Kaczmarek⁸⁵, Beth Y. Karlan¹⁰¹, Karin Kast¹⁰², KConFab Investigators¹⁰³, Sung-Won Kim¹⁰⁴, Irene Konstantopoulou¹⁰⁵, Jacob Korach¹⁰⁶, Yael Laitman², Adriana Lasa¹⁰⁷, Christine Lasset⁴⁰, Conxi Lázaro⁶⁹, Annette Lee¹⁰⁸, Min Hyuk Lee¹⁰⁹, Jenny Lester MPH¹⁰¹, Fabienne Lesueur¹¹⁰, Annelie Liljegren¹¹¹, Noralane M. Lindor¹¹², Michel Longy¹¹³, Jennifer T. Loud¹¹⁴, Karen H. Lu¹¹⁵, Jan Lubinski⁸⁵, Eva Machackova⁷¹, Siranoush Manoukian³¹, Véronique Mari¹¹⁶, Cristina Martínez-Bouzas¹¹⁷, Zoltan Matrai¹¹⁸, Noura Mebirouk¹¹⁰, Hanne E.J. Meijers-Heijboer¹¹⁹, Alfons Meindl⁶⁶, Arjen R. Mensenkamp¹²⁰, Ugnius Mickys¹²¹, Austin Miller¹²², Marco Montagna²³, Kirsten B. Moysich¹²³, Anna Marie Mulligan¹²⁴, Jacob Musinsky¹⁰⁰, Susan L. Neuhausen⁵⁷, Heli Nevanlinna¹²⁵, Joanne Ngeow¹²⁶, Huu Phuc Nguyen¹²⁷, Dieter Niederacher⁹¹, Henriette Roed Nielsen⁹, Finn Cilius Nielsen¹²⁸, Robert L. Nussbaum¹²⁹, Kenneth Offit¹³⁰, Anna Öfverholm¹³¹, Kai-ren Ong¹³², Ana Osorio PhD¹³³, Laura Papi¹³⁴, Janos Papp⁶, Barbara Pasini¹³⁵, Inge Sokilde Pedersen¹³⁶, Ana Peixoto MSc¹³⁷, Nina Peruga MSc⁸⁵, Paolo Peterlongo¹³⁸, Esther Pohl⁸⁶, Nisha Pradhan BA¹⁰⁰, Karolina Prajzendanc⁸⁵, Fabienne Prieur¹³⁹, Pascal Pujol¹⁴⁰, Paolo Radice¹⁴¹, Susan J. Ramus^{142,143}, Johanna Rantala¹⁴⁴, Muhammad Usman Rashid^{3, 145}, Kerstin Rhiem⁸⁶, Mark Robson¹⁴⁶, Gustavo C. Rodriguez¹⁴⁷, Mark T. Rogers¹⁴⁸, Vilius Rudaitis¹⁴⁹, Ane Y. Schmidt¹²⁸, Rita Katharina Schmutzler⁸⁶, Leigha Senter MS¹⁵⁰, Payal D. Shah⁴¹, Priyanka Sharma¹⁵¹, Lucy E. Side¹⁵², Jacques Simard¹⁵³, Christian F. Singer³⁰, Anne-Bine Skytte⁹⁸, Thomas P. Slavin¹⁰, Katie Snape¹⁵⁴, Hagay Sobol¹⁵⁵, Melissa Southey¹⁵⁵, Linda Steele⁵⁷, Doris Steinemann¹⁵⁷, Grzegorz Sukiennicki⁸⁵, Christian Sutter¹⁵⁸, Csilla I. Szabo¹⁵⁹, Yen Y. Tan³⁶, Manuel R. Teixeira¹³⁷, Mary Beth Terry¹⁶⁰, Alex Teulé¹⁶¹, Abigail Thomas MPH¹⁶², Darcy L. Thull MS¹⁶³, Marc Tischkowitz¹⁶⁴, Silvia Tognazzo²³, Amanda Ewart Toland¹⁶⁵, Sabine Topka¹⁰⁰, Alison H Trainer¹⁶⁶, Nadine Tung¹⁶⁷, Christi J. van Asperen¹⁶⁸, Annemieke H. van der Hout¹⁶⁹, Lizet E. van der Kolk¹⁷⁰, Rob B. van der Luit¹⁷¹, Mattias Van Heetvelde⁵¹, Liliana Varesco¹⁷², Raymonda

Varon-Mateeva¹⁷³, Ana Vega¹⁷⁴, Cynthia Villarreal-Garza¹⁷⁵, Anna von Wachenfeldt¹⁷⁶, Lisa Walker¹⁷⁷, Shan Wang-Gohrke¹⁷⁸, Barbara Wappenschmidt⁸⁵, Bernhard H. F. Weber¹⁷⁹, Drakoulis Yannoukakos¹⁰⁵, Sook-Yee Yoon⁸, Cristina Zanzottera³¹, Jamal Zidan¹⁸⁰, Kristin K. Zorn¹⁸¹, Christina G. Hutten Selkirk¹⁸², Peter J. Hulick¹⁸³, Georgia Chenevix-Trench¹⁹, Amanda B. Spurdle¹⁹, Antonis C. Antoniou¹⁸, Katherine L. Nathanson⁴¹ for the CIMBA Consortium

¹Harvard TH Chan School of Public Health and Dana Farber Cancer Institute, 1101 Dana Building, 450 Brookline Ave, Boston, MA 02215, USA

²The Susanne Levy Gertner Oncogenetics Unit, Institute of Human Genetics, Chaim Sheba Medical Center, Ramat Gan 52621, and the Sackler School of Medicine, Tel-Aviv University, Tel-Aviv, Israel

³Molecular Genetics of Breast Cancer, German Cancer Research Center (DKFZ), Im Neuenheimer Feld 580, 69120 Heidelberg, Germany

⁴5841 South Maryland Avenue, MC 2115 Chicago, IL, USA

⁵The Hong Kong Hereditary Breast Cancer Family Registry, Cancer Genetics Center, Hong Kong Sanatorium and Hospital, Hong Kong

⁶Department of Molecular Genetics, National Institute of Oncology, Budapest, Hungary

⁷INBIOMED, Faculty of Medicine, University of Buenos Aires/CONICET and CEMIC, Department of Clinical Chemistry, Medical Direction, Buenos Aires, Paraguay 2155, C1121ABG, Argentina

⁸Cancer Research Initiatives Foundation, Sime Darby Medical Centre, 1 Jalan SS12/1A, Subang Jaya, 47500, Malaysia

⁹Department of Clinical Genetics, Odense University Hospital, Sonder Boulevard 29, Odense C, Denmark

¹⁰Clinical Cancer Genetics, City of Hope, 1500 East Duarte Road, Duarte, California 91010 USA

¹¹Division of Molecular Pathology, Department of Pathology, Hong Kong Sanatorium & Hospital, 1/F Li Shu Fan Block, 2 Village Road, Happy Valley, Hong Kong

¹²Department of Laboratory Medicine and Pathology, and Health Sciences Research, Mayo Clinic, 200 First Street SW, Rochester, Minnesota, USA

¹³Department of Dermatology, University of Utah School of Medicine, 30 North 1900 East, SOM 4B454, Salt Lake City, UT 84132, USA

¹⁴Molecular Oncology Research Center, Barretos Cancer Hospital, Barretos, São Paulo, Brazil

¹⁵1) Department of Preventive Medicine, Seoul National University College of Medicine; 2) Department of Biomedical Science, Seoul National University Graduate School; 3) Cancer Research Center, Seoul National University, 103 Daehak-ro, Jongno-gu, Seoul, Korea

¹⁶Institute of Human Genetics, Pontificia Universidad Javeriana, Carrera 7, Bogota, 11001000, Colombia

¹⁷Cancer Genetics Laboratory, Department of Genetics, University of Pretoria, Private Bag X323, Arcadia 0007, South Africa

¹⁸Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Strangeways Research Laboratory, Worts Causeway, Cambridge, UK

¹⁹Genetics and Computational Biology Department, QIMR Berghofer Medical Research Institute, Herston Road, Brisbane, QLD 4006, Australia

²⁰Department of Clinical Genetics, Academic Medical Center, P.O. Box 22700, 1100 DE Amsterdam, The Netherlands

²¹City of Hope Clinical Cancer Genomics Community Research Network, 1500 East Duarte Road, Duarte, CA 91010, USA

²²Yorkshire Regional Genetics Service, Chapel Allerton Hospital, Leeds, UK

²³Immunology and Molecular Oncology Unit, Veneto Institute of Oncology IOV - IRCCS, Via Gattamelata 64, Padua, Italy

²⁴Department of Clinical Genetics, Helsinki University Hospital, P.O. BOX 160 (Meilahdentie 2), 00029 HUS, Finland

- ²⁵Hereditary Cancer Clinic, Prince of Wales Hospital, High Street, Randwick, NSW 2031 Australia
- ²⁶Lunenfeld-Tanenbaum Research Institute, Sinai Health System, Toronto, Ontario M5G 1X5, Canada; Department of Molecular Genetics, University of Toronto, Toronto, Ontario
- ²⁷Department of Pathology, hus 9, Landspítali-LSH v/Hringbraut, 101 Reykjavik, Iceland
- ²⁸Department of Gynaecology and Obstetrics, University Hospital of Schleswig-Holstein, Campus Kiel, Christian-Albrechts University Kiel, Germany
- ²⁹Department of Breast Medical Oncology and Clinical Cancer Genetics Program, University Of Texas MD Anderson Cancer Center, 1515 Pressler Street, CBP 5, Houston, TX, USA
- ³⁰Dept of OB/GYN and Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria, Waehringer Guertel 18-20, A 1090 Vienna, Austria
- ³¹Unit of Medical Genetics, Department of Medical Oncology and Hematology, Fondazione IRCCS (Istituto Di Ricovero e Cura a Carattere Scientifico) Istituto Nazionale Tumori (INT), Via Giacomo Venezian 1, 20133 Milan, Italy
- ³²Department of Medical Oncology. University Hospital, Vall d'Hebron, Barcelona, Spain
- ³³Division of Cancer Prevention and Genetics, Istituto Europeo di Oncologia (IEO), via Ripamonti 435, 20141 Milan, Italy
- ³⁴Laboratory of Cell Biology, Department of Pathology, hus 9, Landspítali-LSH v/Hringbraut, 101 Reykjavik, Iceland and BMC (Biomedical Centre), Faculty of Medicine, University of Iceland, Vatnsmyrarvegi 16, 101 Reykjavik, Iceland
- ³⁵Human Genetics Group and Genotyping Unit (CEGEN), Human Cancer Genetics Programme, Spanish National Cancer Research Centre (CNIO), Madrid, Spain. Biomedical Network on Rare Diseases (CIBERER), Madrid, Spain
- ³⁶Dept of OB/GYN, Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria, Waehringer Guertel 18-20, 1090 Vienna, Austria
- ³⁷The Institute of Oncology, Chaim Sheba Medical Center, Ramat Gan 52621, Israel
- ³⁸UCSF Cancer Genetics and Prevention Program, San Francisco, CA 94143-1714
- ³⁹Department of Clinical Genetics, Maastricht University Medical Center, P.O. Box 5800, 6202 AZ Maastricht, The Netherlands
- ⁴⁰Unité de Prévention et d'Epidémiologie Génétique, Centre Léon Bérard, 28 rue Laënnec, Lyon, France
- ⁴¹Department of Medicine, Abramson Cancer Center, Perelman School of Medicine at the University of Pennsylvania, 3400 Civic Center Boulevard, Philadelphia, PA 19104, USA
- ⁴²Department of Clinical Genetics, Royal Devon & Exeter Hospital, Exeter, UK
- ⁴³Service de Génétique, Institut Curie, 26, rue d'Ulm, Paris Cedex 05, France
- ⁴⁴Department of Medicine, Huntsman Cancer Institute, 2000 Circle of Hope, Salt Lake City, UT 84112, USA
- ⁴⁵Molecular Oncology Laboratory, Hospital Clinico San Carlos, IdISSC, CIBERONC. Martin Lagos s/n, Madrid, Spain
- ⁴⁶Institute of Human Genetics, University Hospital of Schleswig-Holstein, Campus Kiel, Christian-Albrechts University Kiel, Germany
- ⁴⁷Section of Genetic Oncology, Dept. of Laboratory Medicine, University and University Hospital of Pisa, Pisa, Italy
- ⁴⁸Research Division, Peter MacCallum Cancer Centre, 305 Gratten Street, Melbourne, VIC 3000, Australia
- ⁴⁹CRCHU de Quebec-oncologie, Centre des maladies du sein Deschênes-Fabia, Hôpital du Saint-Sacrement, 1050, chemin Sainte-Foy, Québec Canada
- ⁵⁰Departments of Pediatrics and Medicine, 1150 St. Nicholas Avenue, Columbia University, New York, NY, 10032 USA
- ⁵¹Center for Medical Genetics, Ghent University, De Pintelaan 185, 9000 Gent, Belgium
- ⁵²Department of Clinical Genetics, Family Cancer Clinic, Erasmus University Medical Center, P.O. Box 2040, 3000 CA Rotterdam, The Netherlands

- ⁵³Sheffield Clinical Genetics Service, Sheffield Children's Hospital, Sheffield, UK
- ⁵⁴Department of Clinical Genetics, South Glasgow University Hospitals, Glasgow, UK
- ⁵⁵Unité d'oncogénétique, ICO-Centre René Gauducheau, Boulevard Jacques Monod, 44805 Nantes Saint Herblain Cedex, France
- ⁵⁶Oncogenetics Group, Vall d'Hebron Institute of Oncology (VHIO), Clinical and Molecular Genetics Area, Vall d'Hebron University Hospital, Passeig Vall d'Hebron 119-129, Barcelona, Spain
- ⁵⁷Department of Population Sciences, Beckman Research Institute of City of Hope, Duarte, CA USA
- ⁵⁸Department of Gynaecology and Obstetrics, Ludwig-Maximilian University Munich, Germany
- ⁵⁹Cáncer Hereditario, Instituto de Biología y Genética Molecular, IBGM, Universidad de Valladolid, Centro Superior de Investigaciones Científicas, UVA-CSIC. Valladolid, Spain
- ⁶⁰Institute of Human Genetics, University of Münster, Münster, Germany
- ⁶¹Nottingham Clinical Genetics Service, Nottingham University Hospitals NHS Trust, Nottingham, UK
- ⁶²Oncogenetics Team, The Institute of Cancer Research and Royal Marsden NHS Foundation Trust, Sutton, UK
- ⁶³Department of Clinical Genetics, Lund University Hospital, Lund, Sweden
- ⁶⁴Department of Oncology, Rigshospitalet, Copenhagen University Hospital, Blegdamsvej 9, DK-2100 Copenhagen, Denmark
- ⁶⁵Institute for Medical Informatics, Statistics and Epidemiology, University of Leipzig, Germany
- ⁶⁶Department of Gynaecology and Obstetrics, Division of Tumor Genetics, Klinikum rechts der Isar, Technical University Munich, Germany
- ⁶⁷Genomic Medicine, Manchester Academic Health Sciences Centre, Division of Evolution and Genomic Sciences, University of Manchester, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK
- ⁶⁸Centre de Lutte Contre le Cancer Georges François Leclerc, 1 rue Professeur Marion, BP 77 980, Dijon Cedex, France and Genomic and Immunotherapy Medical Institute, Dijon University Hospital, Dijon, France
- ⁶⁹Molecular Diagnostic Unit, Hereditary Cancer Program, ICO-IDIBELL (Catalan Institute of Oncology-Bellvitge Biomedical Research Institute), CIBERONC, Gran Via de l'Hospitalet, 199-203. 08908 L'Hospitalet. Barcelona, Spain
- ⁷⁰Laboratoire de Génétique Chromosomique, Hôtel Dieu Centre Hospitalier, BP 1125 Chambéry, France
- ⁷¹Department of Cancer Epidemiology and Genetics, Masaryk Memorial Cancer Institute, Zluty kopec 7, Brno, 65653, Czech Republic
- ⁷²Ohio State University /Columbus Cancer Council, Columbus, OH 43221, USA
- ⁷³Oncogenetics Department, Barretos Cancer Hospital, Barretos, São Paulo, Brazil
- ⁷⁴UCLA Schools of Medicine and Public Health, Division of Cancer Prevention & Control Research, Jonsson Comprehensive Cancer Center, 650 Charles Young Drive South, Room A2-125 HS, Los Angeles, CA 90095-6900, USA
- ⁷⁵Cancer Risk and Prevention Clinic, Dana-Farber Cancer Institute, 450 Brookline Avenue, Boston, MA, USA
- ⁷⁶Centre of Familial Breast and Ovarian Cancer, Department of Medical Genetics, Institute of Human Genetics, University Würzburg, Germany
- ⁷⁷Institut Curie, Department of Tumour Biology, Paris, France; Institut Curie, INSERM U830, Paris, France
- ⁷⁸Department of Clinical Genetics, Rigshospitalet 4062, Blegdamsvej 9, København Ø, Denmark
- ⁷⁹Service Régional Oncogénétique Poitou-Charentes, Centre Hospitalier, 79021 Niort
- ⁸⁰Department of Molecular Medicine, University La Sapienza, and Istituto Pasteur - Fondazione Cenci-Bolognetti, viale Regina Elena 291, 00161 Rome, Italy
- ⁸¹Bâtiment Cheney D, Centre Léon Bérard, 28 rue Laënnec, Lyon, France
- ⁸²Ontario Cancer Genetics Network: Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, Toronto, Ontario M5G 1X5, Canada

- ⁸³Department of Pathology and Laboratory Medicine, 3901 Rainbow Boulevard, 4019 Wahl Hall East, MS 3040, University of Kansas Medical Center, Kansas City, Kansas, USA
- ⁸⁴Clinical Genetics Branch, DCEG, NCI, NIH, 9609 Medical Center Drive, Room 6E-454, Bethesda, MD, USA
- ⁸⁵Department of Genetics and Pathology, Pomeranian Medical University, Unii Lubelskiej 1, Szczecin, Poland
- ⁸⁶Center for Familial Breast and Ovarian Cancer, Center for Integrated Oncology (CIO), Medical Faculty, University Hospital Cologne, Cologne, Germany
- ⁸⁷The Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON), Coordinating center: Netherlands Cancer Institute, Amsterdam, The Netherlands
- ⁸⁸Institute of Genetic Medicine, Centre for Life, Newcastle Upon Tyne Hospitals NHS Trust, Newcastle upon Tyne, UK
- ⁸⁹Institute of Human Genetics, University Leipzig, 04107 Leipzig, Germany
- ⁹⁰Family Cancer Clinic, Netherlands Cancer Institute, P.O. Box 90203, 1006 BE Amsterdam, The Netherlands
- ⁹¹Department of Gynaecology and Obstetrics, University Hospital Düsseldorf, Heinrich-Heine University Düsseldorf, Germany
- ⁹²N.N. Petrov Institute of Oncology, St.-Petersburg 197758, Russia
- ⁹³Lombardi Comprehensive Cancer Center, Georgetown University, 3800 Reservoir Road NW, Washington, DC, USA
- ⁹⁴Clinical Genetics, Guy's and St. Thomas' NHS Foundation Trust, London, UK
- ⁹⁵Genetic Counseling Unit, Hereditary Cancer Program, IDIBGI (Institut d'Investigació Biomèdica de Girona), Catalan Institute of Oncology, CIBERONC, Av. França s/n. 1707 Girona, Spain
- ⁹⁶Parkville Familial Cancer Centre, Peter MacCallum Cancer Centre, 305 Gratten Street, Melbourne, VIC 3000, Australia
- ⁹⁷Vilnius University Hospital Santariskiu Clinics, Hereditary Cancer Competence Center Hematology, Oncology and Transfusion Medicine Center Room P519 Santariskiu st. 2, LT-08661 Vilnius, Lithuania
- ⁹⁸Department of Clinical Genetics, Aarhus University Hospital, Brendstrupgaardsvej 21C, Aarhus N, Denmark
- ⁹⁹Department of Epidemiology, Cancer Prevention Institute of California, 2201 Walnut Avenue, Suite 300, Fremont, CA 94538, USA and Department of Health Research and Policy (Epidemiology) and Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA
- ¹⁰⁰Clinical Genetics Research Laboratory, Dept. of Medicine, Memorial Sloan-Kettering Cancer Center, 1275 York Avenue, New York, NY 10044, USA
- ¹⁰¹Women's Cancer Program at the Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, 8700 Beverly Boulevard, Suite 290W, Los Angeles, CA, USA
- ¹⁰²Department of Gynecology and Obstetrics, Medical Faculty and University Hospital Carl Gustav Carus, Technische Universität Dresden, Germany
- ¹⁰³Research Department, Peter MacCallum Cancer Centre, Melbourne, Victoria, Australia and The Sir Peter MacCallum Department of Oncology University of Melbourne, Parkville, Australia
- ¹⁰⁴Department of Surgery, Daerim St. Mary's Hospital, 657 Siheung-daero, Yeongdeungpo-gu, Seoul, Korea
- ¹⁰⁵Molecular Diagnostics Laboratory, INRASTES (Institute of Nuclear and Radiological Sciences and Technology), National Centre for Scientific Research "Demokritos", Patriarchou Gregoriou & Neapoleos str., Aghia Paraskevi Attikis, Athens, Greece
- ¹⁰⁶The Gyneco-Oncology Department, Chaim Sheba Medical Center, Ramat Gan 52621, Israel
- ¹⁰⁷Servicio de Genética-CIBERER U705, Hospital de la Santa Creu i Sant Pau, Barcelona
- ¹⁰⁸The Feinstein Institute for Medical Research 350 Community Drive Manhasset NY
- ¹⁰⁹Department of Surgery, Soonchunhyang University and Seoul Hospital, 59 Daesagwan-Ro, Yongsan-Gu, Seoul, Korea

- ¹¹⁰Institut Curie, PSL Research University, Mines ParisTech, Inserm U900, 26 rue d'Ulm, F-75005 Paris, France
- ¹¹¹Department of Oncology Radiumhemmet and Institution of Oncology and Patology Karolinska University Hospital and Karolinska Institutet
- ¹¹²Department of Health Sciences Research, Mayo Clinic, 13400 E. Scottsdale Blvd., Scottsdale, AZ, USA
- ¹¹³Oncogénétique, Institut Bergonié, 229 cours de l'Argonne, 33076 Bordeaux, France
- ¹¹⁴Clinical Genetics Branch, DCEG, NCI, NIH, 9609 Medical Center Drive, Room 6E-536, Bethesda, MD, USA
- ¹¹⁵Department of Gynecological Oncology and Clinical Cancer Genetics Program, University Of Texas MD Anderson Cancer Center, 1515 Pressler Street, CPB 6, Houston, TX, USA
- ¹¹⁶Centre Antoine Lacassagne, 33 Avenue de Valombrose, Nice, France
- ¹¹⁷Laboratorio de Genética Molecular, Servicio de Genética, Hospital Universitario Cruces, BioCruces Health Research Institute, Spain
- ¹¹⁸Department of Surgery, National Institute of Oncology, Budapest, Hungary
- ¹¹⁹Department of Clinical Genetics, VU University Medical Center, P.O. Box 7057, 1007 MB Amsterdam, The Netherlands
- ¹²⁰Department of Human Genetics, Radboud University Medical Center, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands
- ¹²¹Vilnius university Santariskiu hospital, National Center of Pathology, Baublio st. 5, Vilnius, Lithuania
- ¹²²NRG Oncology, Statistics and Data Management Center, Roswell Park Cancer Institute, Elm St & Carlton St, Buffalo, NY 14263, USA
- ¹²³Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, NY, USA
- ¹²⁴Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, Canada
- ¹²⁵Department of Obstetrics and Gynecology, University of Helsinki and Helsinki University Hospital, Biomedicum Helsinki, P.O. BOX 700 (Haartmaninkatu 8), 00029 HUS, Finland
- ¹²⁶Cancer Genetics Service, Division of Medical Oncology, National Cancer Centre Singapore, 11 Hospital Drive, Singapore 169610
- ¹²⁷Institute of Medical Genetics and Applied Genomics, University of Tuebingen, Germany
- ¹²⁸Center for Genomic Medicine, Rigshospitalet, University of Copenhagen, Denmark
- ¹²⁹513 Parnassus Ave., HSE 901E, San Francisco, CA. 94143 - 0794, USA
- ¹³⁰Clinical Genetics Research Laboratory, Dept. of Medicine, Cancer Biology and Genetics, Memorial Sloan-Kettering Cancer Center, 1275 York Avenue, New York, NY 10044, USA
- ¹³¹Department of Clinical Genetics, Sahlgrenska University Hospital, Gothenburg, Sweden
- ¹³²West Midlands Regional Genetics Service, Birmingham Women's Hospital Healthcare NHS Trust, Edgbaston, Birmingham, UK
- ¹³³Human Genetics Group, Human Cancer Genetics Programme, Spanish National Cancer Research Centre (CNIO), Madrid, Spain. Biomedical Network on Rare Diseases (CIBERER), Madrid, Spain
- ¹³⁴Unit of Medical Genetics, Department of Biomedical, Experimental and Clinical Sciences, University of Florence, Viale Morgagni 50, 50134 Florence, Italy
- ¹³⁵Department of Medical Sciences, University of Turin, Via Santena 19, 10126 Turin, Italy
- ¹³⁶Section of Molecular Diagnostics, Department of Biochemistry, Aalborg University Hospital, Reberbansgade 15, Aalborg, Denmark
- ¹³⁷Department of Genetics, Portuguese Oncology Institute of Porto (IPO Porto), Porto, Portugal, and Biomedical Sciences Institute (ICBAS), University of Porto, Porto, Portugal
- ¹³⁸IFOM, The FIRC (Italian Foundation for Cancer Research) Institute of Molecular Oncology, via Adamello 16, 20139 Milan, Italy.
- ¹³⁹Service de Génétique Clinique Chromosomique et Moléculaire, Hôpital Nord, CHU Saint Etienne, St Etienne cedex 2, France
- ¹⁴⁰Unité d'Oncogénétique, CHU Arnaud de Villeneuve, 34295 Montpellier Cedex 5, France

- ¹⁴¹Unit of Molecular Bases of Genetic Risk and Genetic Testing, Department of Research, Fondazione IRCCS (Istituto Di Ricovero e Cura a Carattere Scientifico) Istituto Nazionale Tumori (INT), c/o Amaedeolab, via GA Amadeo 42, 20133 Milan, Italy
- ¹⁴²School of Women's and Children's Health, UNSW Sydney, Australia
- ¹⁴³The Kinghorn Cancer Centre, Garvan Institute of Medical Research, Australia
- ¹⁴⁴Department of Clinical Genetics, Karolinska University Hospital L5:03, Stockholm S-171 76, Sweden
- ¹⁴⁵Department of Basic Sciences, Shaukat Khanum Memorial Cancer Hospital and Research Centre (SKMCH & RC) 7A, Block R3, Johar Town, Lahore, Punjab 54000, Pakistan
- ¹⁴⁶Clinical Genetics Services, Dept. of Medicine, Memorial Sloan-Kettering Cancer Center, 1275 York Avenue, New York, NY, USA
- ¹⁴⁷Division of Gynecologic Oncology, North Shore University Health System, Clinical Professor, University of Chicago, 2650 Ridge Avenue, Suite 1507 Walgreens, Evanston, IL 60201, USA
- ¹⁴⁸All Wales Medical Genetics Services, University Hospital of Wales, Cardiff, UK
- ¹⁴⁹Vilnius University Hospital Santariskiu Clinics, Centre of Woman's Health and pathology, Department of Gynecology, Santariskiu st. 2, Vilnius, Lithuania
- ¹⁵⁰Clinical Cancer Genetics Program, Division of Human Genetics, Department of Internal Medicine, The Comprehensive Cancer Center, The Ohio State University, Columbus, USA
- ¹⁵¹Department of Hematology and Oncology, University of Kansas Medical Center, Suite 210, 2330 Shawnee Mission Parkway, Westwood, KS, USA
- ¹⁵²North East Thames Regional Genetics Service, Great Ormond Street Hospital for Children NHS Trust, London, UK
- ¹⁵³Genomics Center, Centre Hospitalier Universitaire de Québec Research Center and Laval University, 2705 Laurier Boulevard, Quebec City (Quebec), Canada
- ¹⁵⁴Medical Genetics Unit, St George's, University of London, UK
- ¹⁵⁵Département Oncologie Génétique, Prévention et Dépistage, Institut Paoli-Calmettes, 232 boulevard Sainte-Marguerite, Marseille, France
- ¹⁵⁶Genetic Epidemiology Laboratory, Department of Pathology, University of Melbourne, Parkville, Victoria, Australia
- ¹⁵⁷Institute of Cell and Molecular Pathology, Hannover Medical School, Hannover, Germany
- ¹⁵⁸Department of Human Genetics, University Hospital Heidelberg, Germany
- ¹⁵⁹National Human Genome Research Institute, National Institutes of Health Building 50, Room 5312, 50 South Drive, MSC 004, Bethesda, MD, USA
- ¹⁶⁰Department of Epidemiology, Columbia University, New York, NY, USA
- ¹⁶¹Genetic Counseling Unit, Hereditary Cancer Program, IDIBELL (Bellvitge Biomedical Research Institute), Catalan Institute of Oncology, CIBERONC, Gran Via de l'Hospitalet, 199-203. 08908 L'Hospitalet, Barcelona, Spain
- ¹⁶²Department of Health Sciences Research, Mayo Clinic, 200 First Street SW, Rochester, Minnesota, USA
- ¹⁶³Department of Medicine, Magee-Womens Hospital, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA
- ¹⁶⁴Program in Cancer Genetics, Departments of Human Genetics and Oncology, McGill University, Montreal, Quebec, Canada
- ¹⁶⁵Division of Human Genetics, Departments of Internal Medicine and Cancer Biology and Genetics, Comprehensive Cancer Center, The Ohio State University, 460 W. 12th Avenue, Columbus, OH, USA
- ¹⁶⁶Parkville Familial Cancer Centre, Royal Melbourne Hospital, Melbourne, Australia
- ¹⁶⁷Department of Medical Oncology, Beth Israel Deaconess Medical Center, 330 Brookline Avenue Boston, Massachusetts 02215, USA
- ¹⁶⁸Department of Clinical Genetics, Leiden University Medical Center, P.O. Box 9600, 2300 RC Leiden, The Netherlands
- ¹⁶⁹Department of Genetics, University Medical Center Groningen, University Groningen, The Netherlands

- ¹⁷⁰Family Cancer Clinic, Netherlands Cancer Institute, Amsterdam, The Netherlands
- ¹⁷¹Department of Medical Genetics, University Medical Center Utrecht, The Netherlands
- ¹⁷²Unit of Hereditary Cancer, Department of Epidemiology, Prevention and Special Functions, IRCCS (Istituto Di Ricovero e Cura a Carattere Scientifico) AOU San Martino - IST Istituto Nazionale per la Ricerca sul Cancro, largo Rosanna Benzi 10, 16132 Genoa, Italy
- ¹⁷³Institute of Human Genetics, Campus Virchow Klinikum, Charite Berlin, Germany
- ¹⁷⁴Fundación Pública Galega Medicina Xenómica, calle Choupana s/n, Edificio de Consultas, Planta menos dos Santiago de Compostal, A Coruña, Spain
- ¹⁷⁵Departamento de Investigacion y de Tumores Mamarios del Instituto Nacional de Cancerologia, Mexico City; and Centro de Cancer de Mama del Hospital Zambrano Hellion, Tecnologico de Monterrey, San Pedro Garza Garcia, Nuevo Leon.
- ¹⁷⁶Department of Oncology, Karolinska University Hospital, Stockholm, Sweden
- ¹⁷⁷Oxford Regional Genetics Service, Churchill Hospital, Oxford, UK
- ¹⁷⁸Department of Gynaecology and Obstetrics, University Hospital Ulm, Germany
- ¹⁷⁹Institute of Human Genetics, University Regensburg, Germany
- ¹⁸⁰Institute of Oncology, Rivka Ziv Medical Center, 13000 Zefat, Israel
- ¹⁸¹Magee-Womens Hospital, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA
- ¹⁸²Center for Medical Genetics, NorthShore University HealthSystem, 1000 Central St, Suite 620, Evanston, IL, USA
- ¹⁸³Medical Director, Center for Medical Genetics, North Shore University Health System, Clinical Assistant Professor of Medicine, University of Chicago Pritzker School of Medicine, 1000 Central Street, Suite 620, Evanston, IL 60201, USA

Corresponding Author:

Timothy R. Rebbeck, PhD

1101 Dana, 450 Brookline Avenue

Dana Farber Cancer Institute, Boston, MA 02215

T: 617-632-6128

Email: Timothy_Rebbeck@dfci.harvard.edu

Key Words: *BRCA1*, *BRCA2*, breast cancer, ovarian cancer, mutation, ethnicity, geography

United States NIH, including NCI, funding supported the research presented in this manuscript.

Abstract

The prevalence and spectrum of germline mutations in *BRCA1* and *BRCA2* have been reported in single populations, with the majority of reports focused on Caucasians in Europe and North America. The Consortium of Investigators of Modifiers of *BRCA1/2* (CIMBA) has assembled data on 18,435 families with *BRCA1* mutations and 11,351 families with *BRCA2* mutations ascertained from 69 centers in 49 countries on 6 continents. This study comprehensively describes the characteristics of the 1,650 unique *BRCA1* and 1,731 unique *BRCA2* deleterious (disease-associated) mutations identified in the CIMBA database. We observed substantial variation in mutation type and frequency by geographical region and race/ethnicity. In addition to known founder mutations, mutations of relatively high frequency were identified in specific racial/ethnic or geographic groups that may reflect founder mutations and which could be used in targeted (panel) first pass genotyping for specific populations. Knowledge of the population-specific mutational spectrum in *BRCA1* and *BRCA2* could inform efficient strategies for genetic testing and may justify a more broad-based oncogenetic testing in some populations.

BACKGROUND

Women who carry germline mutations in either *BRCA1* [OMIM 113705] or *BRCA2* [600185] are at a greatly increased risk of breast and ovarian cancers. Estimates of cancer risk associated with *BRCA1* and *BRCA2* mutations vary depending on the population studied. For mutations in *BRCA1*, the estimated average risk of breast and ovarian cancers ranges from 57-65% and 20-50%, respectively (Chen and Parmigiani, 2007; Kuchenbaecker, et al., 2017). For *BRCA2*, average risk estimates range from 35-57% and 5-23%, respectively (Chen and Parmigiani, 2007; Kuchenbaecker, et al., 2017). Mutation-specific cancer risks have been reported that suggest breast cancer cluster regions (BCCR) and ovarian cancer cluster regions (OCCR) exist in both *BRCA1* and *BRCA2* (Kuchenbaecker, et al., 2017; Rebbeck, et al., 2015). The identification of mutations in *BRCA1* or *BRCA2* has important clinical implications, as knowledge of their presence is important for risk assessment and informs medical management for patients. Interventions, such as risk-reducing bilateral mastectomy and salpingo-oophorectomy or annual breast MRI screening, are available to women who carry deleterious *BRCA1* or *BRCA2* mutations to enable early detection of breast cancer and for active risk reduction by risk-reducing surgery (Domchek, et al., 2010; Rebbeck, et al., 2002; Saslow, et al., 2007). The presence of *BRCA1* or *BRCA2* mutations also can influence cancer treatment decisions, principally around the use of platinum agents or poly (ADP-ribose) polymerase (PARP) inhibitors (Lord and Ashworth, 2017) or contralateral risk-reducing mastectomy. Increasing numbers of women are having clinical genetic testing for *BRCA1* and *BRCA2* mutations, and recommendations continue to expand to whom testing should be offered (NCCN, 2017).

In whites drawn from the general populations in North America and the United Kingdom, the prevalence of *BRCA1* and *BRCA2* mutations has been estimated around a broad range from 0.1-0.3%, and 0.1-0.7%, respectively (Peto, et al., 1999; Struewing, et al., 1997; Whittemore, et al., 2004). The Australian Lifepool study, studying a control population consisting of cancer-free women ascertained via population-based mammographic screening program, estimated the overall frequency of *BRCA1* and

BRCA2 mutations to be 0.65% (1:153), with *BRCA1* mutations at 0.20% (1:500) and *BRCA2* mutations at 0.45% (1:222) (Thompson, et al., 2016). Estimates from the Exome Aggregation Consortium (ExAC) are similar, with frequencies of *BRCA1* and *BRCA2* mutations (excluding The Cancer Genome Atlas (TCGA) data) at 0.21% (1:480) and 0.31% (1:327), respectively; or combined at 0.51% (1:195) (Maxwell, et al., 2016). As they do not include large genomic rearrangements, some newer population-based estimates may still under-represent the total number of *BRCA1* and *BRCA2* mutations. Although the overall prevalence of *BRCA1* and *BRCA2* mutations in most general populations is low, many hundreds of thousands of yet-to-be-tested individuals worldwide carry these mutations.

The prevalence of founder mutations in some racial/ethnic groups is much higher. For example, the mutations *BRCA1* c.5266dup (5382insC), *BRCA1* c.68_69del (185delAG) and *BRCA2* c.5946del (6174delT), have a combined prevalence of 2-3% in U.S. Ashkenazi Jews (Roa, et al., 1996; Struwing, et al., 1997; Whittemore, et al., 2004). For these mutations, double heterozygotes in *BRCA1* and *BRCA2* also have been reported (Friedman, et al., 1998; Moslehi, et al., 2000; Ramus, et al., 1997a; Rebbeck, et al., 2016). Several other founder mutations have been identified, including the Icelandic founder mutation *BRCA2* c.771_775del (999del5) (Thorlacius, et al., 1996); the French Canadian mutations *BRCA1* c.4327C>T (C4446T), and *BRCA2* c.8537_8538del (8765delAG) (Oros, et al., 2006b; Tonin, et al., 1999; Tonin, et al., 2001); the *BRCA1* mutations c.181T>G, and c.4034delA in Central-Eastern Europe (Gorski, et al., 2000); the *BRCA1* c.548-4185del in Mexico (Villarreal-Garza, et al., 2015b; Weitzel, et al., 2013)(Villarreal-Garza, et al., 2015b; Weitzel, et al., 2013), the *BRCA2* mutation c.9097dup in Hungary (Ramus, et al., 1997b; Van Der Looij, et al., 2000) and others. These mutations represent the majority of mutations observed in these populations and have been confirmed as true founder mutations as they have common ancestral haplotypes (Neuhausen, et al., 1996, 1998; Oros, et al., 2006a). Recurrent mutations have been identified in other populations, but they represent a smaller proportion of all unique *BRCA1* and *BRCA2* mutations, and have not been characterized as true founder mutations. There are multiple recurrent mutations in Scandinavian, Dutch, French, and

Italian populations (Ferla, et al., 2007). Similarly, a number of recurrent mutations specific to non-European populations also have been reported in Hispanic/Mexican, African-American, Middle Eastern, and Asian populations (Bu, et al., 2016; Ferla, et al., 2007; Kurian, 2010; Lang, et al., 2017; Ossa and Torres, 2016; Villarreal-Garza, et al., 2015b).

The mutational spectra in *BRCA1* and *BRCA2* are best delineated in whites from Europe and North America. However, data on mutational spectra in non-white populations of Asian, African, Mediterranean, South-American and Mexican Hispanic descent have also been reported (Abugattas, et al., 2015; Ahn, et al., 2007; Alemar, et al., 2016; Bu, et al., 2016; Eachkoti, et al., 2007; Ferla, et al., 2007; Gao, et al., 2000; Gonzalez-Hormazabal, et al.; Ho, et al., 2000; Jara, et al., 2006; John, et al., 2007; Kurian, 2010; Laitman, et al.; Lang, et al., 2017; Lee, et al., 2003; Li, et al., 2006; Nanda, et al., 2005; Ossa and Torres, 2016; Pal, et al., 2004; Rodríguez, et al., 2012; Seong, et al., 2009; Sharifah, et al.; Solano, et al., 2017; Song, et al., 2005; Song, et al., 2006; Toh, et al., 2008; Torres, et al., 2007; Troudi, et al., 2007; Villarreal-Garza, et al., 2015b; Vogel, et al., 2007; Weitzel, et al., 2005; Weitzel, et al., 2007; Zhang, et al., 2009). In the current study, we provide a global description of *BRCA1* and *BRCA2* mutations by geography and race/ethnicity from the investigators of the Consortium of Investigators of Modifiers of *BRCA1/2* (CIMBA).

METHODS

Details of centers participating in CIMBA and data collection protocols have been reported previously (Antoniou, et al., 2007). Details of the CIMBA initiative and information about the participating centers can be found at <http://cimba.ccge.medschl.cam.ac.uk/h> (Chenevix-Trench, et al., 2007). All included mutation carriers participated in clinical or research studies at the host institutions after providing informed consent under IRB-approved protocols. Sixty-nine centers and multicenter consortia submitted data that met the CIMBA inclusion criteria (Antoniou, et al., 2007). Only female carriers with pathogenic *BRCA1* and/or *BRCA2* mutations were included in the current analysis. One mutation

carrier per family in the CIMBA database was included in this report. The actual family relationships (e.g., pedigrees) were not available, but a variable that defined family membership supplied by each center was used for this purpose. Less than 1% of families (86 of 29,700) had two family members with two different mutations. In these situations, each mutation observed in the family was included in the analysis. In the case of the 94 dual mutation carriers (i.e., individuals with both *BRCA1* and *BRCA2* mutations), one of the two mutations was chosen at random for inclusion in the analysis.

The CIMBA data set was used to describe the distribution of mutations by effect and function. For the remaining analyses, mutations were excluded if self-reported race/ethnicity data were missing. Pathogenicity of mutation was defined as follows: 1) generating a premature termination codon (PTC), except variants generating a PTC after codon 1854 in *BRCA1* and after codon 3309 of *BRCA2*; 2) large in-frame deletions that span one or more exons; and 3) deletion of transcription regulatory regions (promoter and/or first exon) expected to cause lack of expression of mutant allele. We also included missense variants considered pathogenic by using multifactorial likelihood approaches (Bernstein, et al., 2006; Goldgar, et al., 2004). Mutations that did not meet the above criteria but have been classified as pathogenic by Myriad Genetics, Inc. (Salt Lake City, UT) also were included. Classification of nonsense-mediated decay (NMD) was based on *in-silico* predictions and was not based on molecular classification (Anczukow, et al., 2008).

Contingency table analysis using a chi-square test was used to test for differences in dichotomous variables, as was a t-test for continuous variables. Mutation counts are presented as the number of families with the mutation. Fisher's exact tests were used if sample sizes in any contingency table cell were less than five. Analyses were done in STATA, v. 14.2.

RESULTS

Mutations in *BRCA1* and *BRCA2*

From the 26,861 *BRCA1* and 16,954 *BRCA2* mutation carriers in the CIMBA data set as of June 2017, 18,435 families with *BRCA1* mutations and 11,351 families with *BRCA2* mutations were studied to count only one occurrence of a mutation per family. **Figure 1** shows the countries that contributed mutations to this report. From among these families, 1,650 unique *BRCA1* and 1,731 unique *BRCA2* mutations were identified. The unique mutations and number of families in which each mutation was observed are listed in **Supplementary Table 1**. In each gene, the five most common mutations (including founder mutations) accounted for 33% of all mutations in *BRCA1* (8,739 of 26,861 mutation carriers) and 19% of all mutations in *BRCA2* (3,244 of 16,954 mutation carriers). A web site containing information about the most common mutations reported here can be found at: <http://apps.ccge.medschl.cam.ac.uk/consortia/cimba/>. This information may be periodically updated as new data become available.

Mutation Type and Effect

Table 1 presents a summary of the type of *BRCA1* or *BRCA2* mutations and their predicted effect on transcription and translation. The most common mutation type was frameshift followed by nonsense. The most common effect of *BRCA1* and *BRCA2* mutations was premature translation termination and most of the mutant mRNAs were predicted to undergo nonsense-mediated mRNA decay (NMD) (Anczukow, et al., 2008). Despite having the same spectrum of mutations in *BRCA1* and *BRCA2*, the frequency distribution by mutation type, effect, or function differed significantly ($p < 0.05$) between *BRCA1* and *BRCA2* mutation carriers for many groups, as shown in **Table 1**. These observed differences are largely because genomic rearrangements and missense mutations account for a much higher proportion of mutations in *BRCA1* when compared with *BRCA2*, as previously described (Welch and King, 2001).

We and others have found that breast (BCCR) and ovarian (OCCR) cancer cluster regions exist that may confer differential cancer risks (Gayther, et al., 1997; Gayther, et al., 1995; Kuchenbaecker, et al.,

2017; Rebbeck, et al., 2015). **Figure 2** reports the relative frequency of mutations in the BCCR and OCCR by race/ethnicity. Compared with whites, we observed differences in the relative frequency of mutations in the *BRCA1* BCCR and OCCR in Asians and Hispanics, and in the *BRCA2* OCCR in Hispanics. To the degree that the mutations within the BCCRs and OCCRs conferred differential cancer risks, these data suggest that *BRCA1* and *BRCA2* mutation-associated cancer risks may vary by race/ethnicity.

Geography and Race/Ethnicity

The most common mutations by country are summarized in **Table 2** (*BRCA1*) and **Table 3** (*BRCA2*). The locations of the mutations that were observed in African American, Asian, and Hispanic populations are depicted in **Figure 3** (*BRCA1*) and **Figure 4** (*BRCA2*). Some countries (Albania, Bosnia, Costa Rica, Ireland, Honduras, Japan, Norway, Peru, Philippines, Qatar, Saudi Arabia, Romania, Venezuela and Turkey) contributed fewer than 10 mutation carriers to the CIMBA database. Many of these mutations were submitted to the central database by CIMBA centers that ascertained these patients, but these patients originated from a different country. Based on such small numbers, it was impossible to make inferences about the relative importance of mutations in these locations. A description of the major ethnicity by country is provided in **Supplementary Table 2**.

The mutational distribution among the major racial/ethnic groups and by geography are summarized in **Tables 4** and **5**. Table 4 includes only those individuals for whom self-identified race/ethnicity was recorded. Note that in some countries it is prohibited to collect data on race and ethnicity, so this information is missing. Among the 10 most common *BRCA1* mutations in each racial/ethnic group, a few were seen in several populations, including the recurrent Jewish and Eastern European founder mutations c.5266dup (5382insC) and c.68_69del (185delAG); c.815_824dup in African-Americans and Hispanics; c.3756_3759del in Caucasian and Jews; and c.5503C>T and c.3770_3771del in Asians and Jews. Similarly, recurrent mutations in *BRCA2* included c.5946del (6174delT) in whites and Jews;

c.2808_2811del in whites, African Americans, Asians, Hispanics, and Jews; c.6275_6276del in whites and Hispanics; c.3847_3848del in whites and Jews; c.658_659del in African Americans and Hispanics; and c.3264dup in Hispanics and Jews. The majority of other recurrent *BRCA1* and *BRCA2* mutations were only observed within a single racial/ethnic group, particularly African Americans, Asians, and Hispanics. Of note, the vast majority of women who self-identified as Jewish carry the Ashkenazi Jewish founder mutations *BRCA1* c.5266dup and c.68_69del and *BRCA2* c.5946del. Only 72 (3.9%) of 1,852 *BRCA1* mutation carrier families and 55 (5.6%) of 990 *BRCA2* mutation carrier families who self-identified as being Jewish carried other (non-founder) mutations. However, since many individuals of self-identified Jewish ancestry are only tested for the three founder mutations, this number is likely to be underestimated.

In African Americans, the majority of *BRCA1* mutations were not observed in any other racial/ethnic group, implying these mutations may be of African origin. In Hispanics, the most common *BRCA1* mutations also were observed among individuals from other regions who did not self-identify as Hispanic, including *BRCA1* c.3331_3334del (also observed in Australia, Europe, USA, and the UK), and *BRCA1* c.68_69del (the Jewish founder mutation) (Weitzel, et al., 2013; Weitzel, et al., 2005). The *BRCA1* c.815_824dup mutation has been reported as being of African origin, but has also been reported as a recurrent mutation in Mexican-Americans, perhaps as a reflection of the complex continental admixture of this population (Villarreal-Garza, et al., 2015b). *BRCA1* c.390C>A and c.5496_5506delinsA were most commonly found in the Asian population. In *BRCA2*, c.2808_2811del was found among the 10 most frequent mutations in all races/ethnicities.

Recurrent Mutations

As expected, the most common mutations in the entire data set were the founder mutations *BRCA1* c.5266dup (5382insC), *BRCA1* c.68_69del (185delAG), and *BRCA2* c.5946del (6174delT). In part, the high frequency of these mutations is a consequence of panels that facilitate testing for these three

mutations in women of Jewish descent. However, these two *BRCA1* mutations also are relatively common in regions with a low proportion of individuals who self-identify as Jewish (e.g., Hungary, Czech Republic, France, Germany, Italy, Poland Spain, Russia, and UK). *BRCA1* c.5266dup is a founder mutation thought to have originated 1800 years ago in Scandinavia/Northern Russia, entering the Ashkenazi-Jewish population 400-500 years ago, and thus has origins and a spread pattern independent of the Ashkenazim (Hamel, et al., 2011). Haplotype studies have been used to determine the origin of *BRCA1* c.68_69delAG in populations not considered to have a high proportion of Jewish ancestry. In some populations, such as the Hispanics in the USA and Latin American, it is associated with the Ashkenazi Jewish haplotype, presumably due to unrecognized (Jewish) ancestry (Ah Mew, et al., 2002; Velez, et al., 2012; Weitzel, et al., 2005). In other populations, such as Pakistani and Malaysians, where *BRCA1* c.68_69del is a recurrent mutation, it appears to have arisen independently, as it is carried on a distinct haplotype (Kadalmani, et al., 2007; Rashid, et al., 2006). A different haplotype was also reported for several British families (the 'Yorkshire haplotype') that is distinct from both the Jewish and the Indian-Pakistani haplotypes (Laitman, et al., 2013; Neuhausen, et al., 1996).

The only locations in which these three founder mutations were not commonly observed were Belgium and Iceland. Iceland has another founder mutation (i.e., *BRCA2* c.771_775del). Yet other founder mutations included *BRCA1* c.4327C>T and *BRCA2* c.8537_8538del in Quebec. This latter mutation in *BRCA2* also is the most common mutation in high-risk families in Sardinia (Pisano, et al., 2000) and was also reported in a few Jewish Yemenite families, with a distinct haplotype (Palomba, et al., 2007). The *BRCA1* c.181T>G mutation was observed in Central Europe (Austria, Czech Republic, Germany, Hungary, Italy and Poland), but also observed in the US, Argentina, Latvia, Lithuania and Israel. This mutation has been found on a common haplotype in individuals of Polish and Ashkenazi Jewish ancestry, suggesting it is an Eastern European founder mutation (Kaufman, et al., 2009). The large rearrangement mutation in *BRCA1* c.548-?4185+?del (ex9-12del) appears to be an important founder

mutation in Mexico, with findings of a common haplotype and an estimated age at 74 generations (~1,500 years) (Weitzel, et al., 2013).

We observed a number of other recurrent mutations. *BRCA1* c.3331_3334del comprised more than half of all mutations identified in Colombia, consistent with a previous report that this is a founder mutation in the Colombian population (Torres, et al., 2007). However, this mutation has not been found at high rates in a second Colombian population (Cock-Rada, et al., 2017). *BRCA2* c.2808_2811del was frequently observed, not only as the most common mutation in France and Colombia, but also in other Western and Southern European countries, and destinations to which individuals from these countries have migrated. It estimated to have arisen approximately 80 (46-134) generations ago. However, due to the diversity of the haplotypes, multiple independent origins could not be ruled out (Neuhausen, et al., 1998). *BRCA2* c.6275_6276del was a recurrent *BRCA2* mutation in Australia, the UK, Belgium, Spain, the Netherlands, and North America. This mutation has been estimated to have originated 52 (24-98) generations ago from a single founder (Neuhausen, et al., 1998). Recurrent or founder mutations were observed in diverse populations. For example, the c.115T>G (Cys39Gly) mutation has been described in Greenlanders (Hansen, et al., 2009). The c.2641G >T and c.7934del mutations have both been reported as founder mutation in South African Afrikaners (Reeves, et al., 2004).

DISCUSSION

We have reported worldwide distribution of *BRCA1* and *BRCA2* mutations curated in the CIMBA dataset. These results may aid in the understanding of the mutation distribution in specific populations as well as imparting clinical and biological implications for our understanding of *BRCA1*- and *BRCA2*-associated carcinogenesis.

Clinical testing for *BRCA1* and *BRCA2* mutations has benefited substantially from knowledge about common mutations in specific populations. In many countries, the three Ashkenazi-Jewish founder

mutations are offered as a mutation testing panel for self-reported Ashkenazim, based on their frequency. This approach is much less expensive than comprehensive gene sequencing. The identification of commonly-occurring mutations in other populations could lead to more efficient and cost-effective mutation testing for *BRCA1* and *BRCA2*. For example, Villareal-Garza et al. (Villarreal-Garza, et al., 2015a) have developed the HISPANEL of mutations that optimizes testing in Hispanic/Latino populations. In the present study, we have identified mutations that may exist at a sufficient prevalence to warrant consideration for population-specific mutation testing panels. Criteria for developing such panels for *BRCA1* and *BRCA2* mutation screening are not available. However, mutations that are in a specific population and that capture a sufficient percentage of mutations in high risk individuals and families in that population may be appropriate for use in targeted genetic testing. Before such panels can be developed, population-based studies of mutation frequency in specific populations should be undertaken. The data reported herein provide a list of the recurrent mutations around which such panels could be developed, but the frequencies are not population-based, particularly in settings where founder mutations are preferentially screened (e.g., the Jewish founder panels). Similarly, putative founder mutations identified by assessing common ancestral origins of specific mutations (rather than just high prevalence; **Table 5**) may form the basis of population-specific *BRCA1* and *BRCA2* mutation screening panels.

We report the distribution of *BRCA1* and *BRCA2* mutations in nearly 30,000 families of bona-fide disease-associated mutations. The strengths of this report include the large sample size that reflects a geographically and racially/ethnically diverse set of *BRCA1* and *BRCA2* mutation carriers. However, some limitations need to be considered. First, the sample set presented here does not reflect a systematic study of these populations or races/ethnicities; the data reflect patterns of recruitment (e.g., individuals with higher risk or prior diagnosis of cancer who consented to participate in research protocols) that contributed to the CIMBA consortium. Certain racial/ethnic or socio-demographic groups are under- or over-represented or missing in our data set and, as a consequence, mutations may be

over- or under-represented. For example, the existence of a commercial panel of three Jewish founder mutations enhances genetic testing for those mutations. As a result, the most frequently observed mutations in some populations (e.g., the USA) reflect the widespread use of this testing panel in the USA population. Similar arguments may also apply for other populations, where testing for certain founder mutations may be more frequent. Therefore the relative frequencies of mutations by population in the present study may be subject to such testing biases. Comparing the relative frequencies is also complicated by the inclusion of related individuals.

Second, although the CIMBA data represent most regions around the world, there are limitations related to which groups of individuals have been tested and which centers contributed data. In particular, non-white ancestry populations are still under-represented in research reports of mutation spectrum and frequency. Genetic testing in the developing world remains limited.

Third, we presented the mutations in terms of type or effect (**Table 1**), but these designations are not always based on experimental evidence. For example, NMD mutation status is almost always defined by a prediction rule rather than *in vitro* experiments that confirm the presence of nonsense mediated decay.

Fourth, we presented the occurrence of putative founder mutations. Some of these founder mutations (e.g., *BRCA1* c.68_69del, *BRCA2* c.771_775del) have been demonstrated to be true founder mutations based on actual ancestry analyses. Others, however, have only been identified as occurring commonly in certain populations, but haplotype or similar analyses of founder status may not have been done.

Fifth, our analysis was based on self-reported race/ethnicity of study participants, but this information may misclassify some groups of individuals. For example, some Middle Eastern groups may have been classified as “Caucasian” based on the data available, but in fact may represent a distinct group that

was not captured here. Moreover, in some large centers participating in CIMBA, collecting information on race/ethnicity is prohibited and these mutation carriers were excluded from the comparisons.

Finally, we evaluated mutations by racial/ethnic and geographic designations, but some of these may be misclassified. For example, while *BRCA1* c.68_69del has been shown to arise independently of the Jewish founder mutation in Pakistan (Rashid, et al., 2006), we cannot determine if the identified group also contains some Ashkenazi Jewish individuals.

The data presented herein provide new insights into the worldwide distribution of *BRCA1* and *BRCA2* mutations. The identification of recurrent mutations in some racial/ethnic groups or geographical locations raises the possibility of defining more efficient strategies for genetic testing. Three Jewish founder mutations *BRCA1* c.5266dup (5382insC) and *BRCA1* c.68_69del (185delAG) and *BRCA2* c.5946del (6174delT) have long been used as a primary genetic screening test for women of Jewish descent. The identification here of other recurrent mutations in specific populations may similarly provide the basis for other mutation-specific panels. For example, *BRCA1* c.5266dup (5382insC) may be a useful as a single mutation screening test in Central-Eastern European populations before undertaking full sequencing. However, this basic test may be supplemented with screening for *BRCA1* c.181T>G, as the second most common mutation of the region, and for some special cases, to include most common Hungarian *BRCA2* founder mutation c.9097dup (9326insA) for those with Hungarian ancestry (van der Looij, et al., 2000, Ramus, et al., 1997b). In Iceland, only two mutations were reported: the founder mutation *BRCA2* c.771_775del and the rarer *BRCA1* c.5074G>A (Bergthorsson, et al., 1998). A number of other situations can be identified in which specific mutations explain a large proportion of the total mutations observed in a population. These and other such examples suggest that targeted mutation testing panels which include specific mutations could be developed for use in specific populations. Finally, we focused on female *BRCA1* and *BRCA2* mutation carriers in this report. However, the growing knowledge about *BRCA1* and *BRCA2*-associated cancers in men, particularly

prostate cancer (Ostrander and Udler, 2008; Pritchard, et al., 2016), suggests that the information presented herein will also have value in genetic testing of men.

ACKNOWLEDGMENTS

Study	Funding	Acknowledgements
CIMBA	The CIMBA data management and data analysis were supported by Cancer Research – UK grants C12292/A20861, C12292/A11174. ACA is a Cancer Research -UK Senior Cancer Research Fellow. GCT and ABS are NHMRC Research Fellows. iCOGS: the European Community's Seventh Framework Programme under grant agreement n° 223175 (HEALTH-F2-2009-223175) (COGS), Cancer Research UK (C1287/A10118, C1287/A 10710, C12292/A11174, C1281/A12014, C5047/A8384, C5047/A15007, C5047/A10692, C8197/A16565), the National Institutes of Health (CA128978) and Post-Cancer GWAS initiative (1U19 CA148537, 1U19 CA148065 and 1U19 CA148112 - the GAME-ON initiative), the Department of Defence (W81XWH-10-1-0341), the Canadian Institutes of Health Research (CIHR) for the CIHR Team in Familial Risks of Breast Cancer (CRN-87521), and the Ministry of Economic Development, Innovation and Export Trade (PSR-SIIRI-701), Komen Foundation for the Cure, the Breast Cancer Research Foundation, and the Ovarian Cancer Research Fund. The PERSPECTIVE project was supported by the Government of Canada through Genome Canada and the Canadian Institutes of Health Research, the Ministry of Economy, Science and Innovation through Genome Québec, and The Quebec Breast Cancer Foundation.	All the families and clinicians who contribute to the studies; Sue Healey, in particular taking on the task of mutation classification with the late Olga Sinilnikova; Maggie Angelakos, Judi Maskiell, Gillian Dite, Helen Tsimiklis
BCFR - all	This Breast Cancer Family Registry (BCFR) is supported by grant UM1 CA164920 from the USA National Cancer Institute. The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centers in the BCFR, nor does mention of trade names, commercial products, or organizations imply endorsement by the USA Government or the	

Study	Funding	Acknowledgements
	BCFR.	
BCFR-AU		Maggie Angelakos, Judi Maskiell, Gillian Dite, Helen Tsimiklis.
BCFR-NY		We wish to thank members and participants in the New York site of the Breast Cancer Family Registry for their contributions to the study.
BCFR-ON		We wish to thank members and participants in the Ontario Familial Breast Cancer Registry for their contributions to the study.
BFBOCC-LT	BFBOCC is partly supported by: Lithuania (BFBOCC-LT): Research Council of Lithuania grant SEN-18/2015	BFBOCC-LT acknowledge Laimonas Griškevičius. BFBOCC-LV acknowledge Drs Janis Eglitis, Anna Krilova and Aivars Stengrevics.
BIDMC	BIDMC is supported by the Breast Cancer Research Foundation	
BMBSA	BRCA-gene mutations and breast cancer in South African women (BMBSA) was supported by grants from the Cancer Association of South Africa (CANSA) to Elizabeth J. van Rensburg	BMBSA wish to thank the families who contribute to the BMBSA study
BRICOH	SLN was partially supported by the Morris and Horowitz Families Endowed Professorship.	
CEMIC	This work is funded by CONICET and Instituto Nacional del Cancer, Ministerio de Salud de la Nacion Argentina (1995/15)	We thank to Florencia Cardoso, Natalia Liria and Pablo Mele in their biospecimen and data management.
CNIO	CNIO study is partially funded by the Spanish Ministry of Health PI16/00440 supported by FEDER funds, the Spanish Ministry of Economy and Competitiveness (MINECO) SAF2014-57680-R and the Spanish Research Network on Rare diseases (CIBERER)	We thank Alicia Barroso, Rosario Alonso and Guillermo Pita for their assistance.

Study	Funding	Acknowledgements
COH-CCGCRN	<p>City of Hope Clinical Cancer Genomics Community Network and the Hereditary Cancer Research Registry, supported in part by the Breast Cancer Research Foundation, by Award Number RC4CA153828 (PI: J. Weitzel) from the National Cancer Institute and the Office of the Director, National Institutes of Health, and by the National Cancer Institute of the National Institutes of Health under Award Number R25CA171998 (PIs: K. Blazer and J. Weitzel). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health</p>	
CONSIT TEAM	<p>Associazione Italiana Ricerca sul Cancro (AIRC; IG2014 no.15547) to P. Radice; Funds from Italian citizens who allocated the 5x1000 share of their tax payment in support of the Fondazione IRCCS Istituto Nazionale Tumori, according to Italian laws (INT-Institutional strategic projects '5x1000') to S Manoukian; Associazione Italiana per la Ricerca sul Cancro IG17734; Italian Ministry of University and Research, PRIN projects; Istituto Pasteur-Fondazione Cenci Bolognetti to G. Giannini ; FiorGen Foundation for Pharmacogenomics to L. Papi ; Funds from Italian citizens who allocated the 5x1000 share of their tax payment in support of the IRCCS AOU San Martino - IST according to Italian laws (institutional project) to L. Varesco ; Associazione Italiana Ricerca sul Cancro (AIRC; IG2015 no.16732) to P. Peterlongo</p>	<p>Bernard Peissel, Milena Mariani and Daniela Zaffaroni of the Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy; Davide Bondavalli, Maria Rosaria Calvello and Irene Feroce of the Istituto Europeo di Oncologia, Milan, Italy; Alessandra Viel and Riccardo Dolcetti of the CRO Aviano National Cancer Institute, Aviano (PN), Italy; Francesca Vignolo-Lutati of the University of Turin, Turin, Italy; Gabriele Capone of the University of Florence, Florence, Italy; Laura Ottini of the "Sapienza" University, Rome, Italy; Viviana Gismondi of the IRCCS AOU San Martino – IST, Istituto Nazionale per la Ricerca sul Cancro, Genoa, Italy; Maria Grazia Tibiletti and Daniela Furlan of the Ospedale di Circolo-Università dell'Insubria, Varese, Italy; Antonella Savarese and Aline Martayan of the Istituto Nazionale Tumori Regina Elena, Rome, Italy; Stefania Tommasi and Brunella Pilato of the Istituto Nazionale Tumori "Giovanni Paolo II" - Bari, Italy, and the personnel of the Cogentech Cancer Genetic Test Laboratory, Milan, Italy.</p>

Study	Funding	Acknowledgements
DFCI	This research has been supported by R01-CA08534 and R01-CA102776 to TRR.	
DEMOKRITOS	This research has been co-financed by the European Union (European Social Fund – ESF) and Greek national funds through the Operational Program "Education and Lifelong Learning" of the National Strategic Reference Framework (NSRF) - Research Funding Program of the General Secretariat for Research & Technology: SYN11_10_19 NBCA. Investing in knowledge society through the European Social Fund.	
DKFZ	The DKFZ study was supported by the DKFZ and in part by the SKMCH & RC, Lahore, Pakistan and the Pontificia Universidad Javeriana, Bogota, Colombia .	We thank all participants, clinicians, family doctors, researchers, and technicians for their contributions and commitment to the DKFZ study and the collaborating groups in Lahore, Pakistan (Noor Muhammad, Sidra Gull, Seerat Bajwa, Faiz Ali Khan, Humaira Naeemi, Saima Faisal, Asif Loya, Mohammed Aasim Yusuf) and Bogota, Colombia (Ignacio Briceno, Fabian Gil).
EMBRACE	EMBRACE is supported by Cancer Research UK Grants C1287/A10118 and C1287/A11990. D. Gareth Evans and Fiona Lalloo are supported by an NIHR grant to the Biomedical Research Centre, Manchester. The Investigators at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust are supported by an NIHR grant to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust. Ros Eeles and Elizabeth Bancroft are supported by Cancer Research UK Grant C5047/A8385. Ros Eeles is also supported by NIHR support to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust	RE is supported by NIHR support to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust
FCCC	The authors acknowledge support from The University of Kansas Cancer Center (P30 CA168524) and the Kansas Bioscience Authority Eminent Scholar Program. A.K.G. was funded by 5U01CA113916, R01CA140323, and by the Chancellors Distinguished Chair in Biomedical Sciences Professorship.	We thank Ms. JoEllen Weaver and Dr. Betsy Bove for their technical support.

Study	Funding	Acknowledgements
FPGMX	This work was partially supported by FISPI05/2275 and Mutua Madrileña Foundation (FMMA).	We would like to thank Marta Santamariña, Ana Blanco, Miguel Aguado, Uxía Esperón and Belinda Rodríguez for their contribution with the study.
GC-HBOC	The German Consortium of Hereditary Breast and Ovarian Cancer (GC-HBOC) is supported by the German Cancer Aid (grant no 110837), Rita K. Schmutzler.	The Regensburg HBOC thanks Dr. Ivana Holzhauser and Dr. Ines Schönbuchner for their contributions to the study
GEMO	The study was supported by the Ligue Nationale Contre le Cancer; the Association “Le cancer du sein, parlons-en!” Award; the Canadian Institutes of Health Research for the "CIHR Team in Familial Risks of Breast Cancer" program and the French National Institute of Cancer (INCa).	Genetic Modifiers of Cancer Risk in <i>BRCA1</i> or <i>BRCA2</i> Mutation Carriers (GEMO) study : National Cancer Genetics Network «UNICANCER Genetic Group», France. We wish to pay a tribute to Olga M. Sinilnikova, who with Dominique Stoppa-Lyonnet initiated and coordinated GEMO until she sadly passed away on the 30th June 2014, and to thank all the GEMO collaborating groups for their contribution to this study. GEMO Collaborating Centers are: Coordinating Centres, Unité Mixte de Génétique Constitutionnelle des Cancers Fréquents, Hospices Civils de Lyon - Centre Léon Bérard, & Equipe «Génétique du cancer du sein», Centre de Recherche en Cancérologie de Lyon: Olga Sinilnikova†, Sylvie Mazoyer, Francesca Damiola, Laure Barjhoux, Carole Verny-Pierre, Mélanie Léone, Nadia Boutry-Kryza, Alain Calender, Sophie Giraud; and Service de Génétique Oncologique, Institut Curie, Paris: Claude Houdayer, Etienne Rouleau, Lisa Golmard, Agnès Collet, Virginie Moncoutier, Muriel Belotti, Camille Elan, Catherine Nogues, Emmanuelle Fourme, Anne-Marie Birot. Institut Gustave Roussy, Villejuif: Brigitte Bressac-de-Paillerets,

Study	Funding	Acknowledgements
		<p>Olivier Caron, Marine Guillaud-Bataille. Centre Jean Perrin, Clermont–Ferrand: Yves-Jean Bignon, Nancy Uhrhammer. Centre Léon Bérard, Lyon: Christine Lasset, Valérie Bonadona, Sandrine Handallou. Centre François Baclesse, Caen: Agnès Hardouin, Pascaline Berthet, Dominique Vaur, Laurent Castera. Institut Paoli Calmettes, Marseille: Hagay Sobol, Violaine Bourdon, Tetsuro Noguchi, Audrey Remenieras, François Eisinger. CHU Arnaud-de-Villeneuve, Montpellier: Isabelle Coupier, Pascal Pujol. Centre Oscar Lambret, Lille: Jean-Philippe Peyrat, Joëlle Fournier, Françoise Révillion, Philippe Vennin†, Claude Adenis. Centre Paul Strauss, Strasbourg: Danièle Muller, Jean-Pierre Fricker. Institut Bergonié, Bordeaux: Emmanuelle Barouk-Simonet, Françoise Bonnet, Virginie Bubien, Nicolas Sevenet, Michel Longy. Institut Claudius Regaud, Toulouse: Christine Toulas, Rosine Guimbaud, Laurence Gladieff, Viviane Feillel. CHU Grenoble: Dominique Leroux, Hélène Dreyfus, Christine Rebischung, Magalie Peysselon. CHU Dijon: Fanny Coron, Laurence Faivre. CHU St-Etienne: Fabienne Prieur, Marine Lebrun, Caroline Kientz. Hôtel Dieu Centre Hospitalier, Chambéry: Sandra Fert Ferrer. Centre Antoine Lacassagne, Nice: Marc Frénay. CHU Limoges: Laurence Vénat-Bouvet. CHU Nantes: Capucine Delnatte. CHU Bretonneau, Tours: Isabelle Mortemousque.</p>

Study	Funding	Acknowledgements
		Groupe Hospitalier Pitié-Salpêtrière, Paris: Florence Coulet, Chrystelle Colas, Florent Soubrier, Mathilde Warcoin. CHU Vandoeuvre-les-Nancy: Johanna Sokolowska, Myriam Bronner. CHU Besançon: Marie-Agnès Collonge-Rame, Alexandre Damette. Creighton University, Omaha, USA: Henry T. Lynch, Carrie L. Snyder.
GEORGETOWN	CI received support from the Non-Therapeutic Subject Registry Shared Resource at Georgetown University (NIH/NCI grant P30-CA051008), the Fisher Center for Familial Cancer Research, and Swing Fore the Cure.	
HCBARRETOS	This study was supported by Barretos Cancer Hospital, FINEP - CT-INFRA (02/2010) and FAPESP (2013/24633-2).	We wish to thank members of the Center of Molecular Diagnosis, Oncogenetics Department and Molecular Oncology Research Center of Barretos Cancer Hospital for their contributions to the study.
G-FAST	Bruce Poppe is a senior clinical investigator of FWO. Mattias Van Heetvelde obtained funding from IWT.	We wish to thank the technical support of Ilse Coene en Brecht Crombez.
HCSC	Was supported by a grant RD12/0036/0006 and 15/00059 from ISCIII (Spain), partially supported by European Regional Development FEDER funds	We acknowledge Alicia Tosar and Paula Diaque for their technical assistance
HEBCS	The HEBCS was financially supported by the Helsinki University Hospital Research Fund, Academy of Finland (266528), the Finnish Cancer Society and the Sigrid Juselius Foundation.	HEBCS would like to thank Taru A. Muranen and Johanna Kiiski, Drs. Carl Blomqvist and Kirsimari Aaltonen and RNs Irja Erkkilä and Virpi Palola for their help with the HEBCS data and samples. HEBCS would like to thank Dr. Kristiina Aittomäki, Taru A. Muranen, Drs. Carl Blomqvist and Kirsimari Aaltonen and RNs Irja Erkkilä and Virpi Palola for their help with the HEBCS data and samples.

Study	Funding	Acknowledgements
HEBON	<p>The HEBON study is supported by the Dutch Cancer Society grants NKI1998-1854, NKI2004-3088, NKI2007-3756, the Netherlands Organization of Scientific Research grant NWO 91109024, the Pink Ribbon grants 110005 and 2014-187.WO76, the BBMRI grant NWO 184.021.007/CP46 and the Transcan grant JTC 2012 Cancer 12-054. HEBON thanks the registration teams of Dutch Cancer Registry (IKNL; S. Siesling, J. Verloop) and the Dutch Pathology database (PALGA; L. Overbeek) for part of the data collection.</p>	<p>The Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON) consists of the following Collaborating Centers: Coordinating center: Netherlands Cancer Institute, Amsterdam, NL: M.A. Rookus, F.E. van Leeuwen, S. Verhoef, M.K. Schmidt, N.S. Russell, J.L. de Lange, R.Wijnands; Erasmus Medical Center, Rotterdam, NL: J.M. Collée, A.M.W. van den Ouweland, M.J. Hooning, C. Seynaeve, C.H.M. van Deurzen, I.M. Obdeijn; Leiden University Medical Center, NL: J.T. Wijnen, R.A.E.M. Tollenaar, P. Devilee, T.C.T.E.F. van Cronenburg; Radboud University Nijmegen Medical Center, NL: C.M. Kets; University Medical Center Utrecht, NL: M.G.E.M. Ausems, R.B. van der Luijt, C.C. van der Pol; Amsterdam Medical Center, NL: C.M. Aalfs, T.A.M. van Os; VU University Medical Center, Amsterdam, NL: J.J.P. Gille, Q. Waisfisz; Maastricht University Medical Center: University Hospital Maastricht, NL: E.B. Gómez-Garcia; University Medical Center Groningen, NL: J.C. Oosterwijk, A.H. van der Hout, M.J. Mourits, G.H. de Bock; The Netherlands Foundation for the detection of hereditary tumours, Leiden, NL: H.F. Vasen; The Netherlands Comprehensive Cancer Organization (IKNL): S. Siesling, J. Verloop; The Dutch Pathology Registry (PALGA): L.I.H. Overbeek.</p>

Study	Funding	Acknowledgements
HRBCP	HRBCP is supported by The Hong Kong Hereditary Breast Cancer Family Registry and the Dr. Ellen Li Charitable Foundation, Hong Kong	We wish to thank Hong Kong Sanatorium and Hospital for their continued support
HUNBOCS	Hungarian Breast and Ovarian Cancer Study was supported by Hungarian Research Grants KTIA-OTKA CK-80745, NKFIH/ OTKA K-112228 and the Norwegian EEA Financial Mechanism Hu0115/NA/2008-3/OP-9	We wish to thank the Hungarian Breast and Ovarian Cancer Study Group members (Janos Papp, Tibor Vaszko, Aniko Bozsik, Timea Pocza, Zoltan Matrai, Gabriella Ivady , Judit Franko, Maria Balogh, Gabriella Domokos, Judit Ferenczi, Department of Molecular Genetics , National Institute of Oncology, Budapest, Hungary) and the clinicians and patients for their contributions to this study.
HVH		We wish to thank the Oncogenetics Group (VHIO) and the High Risk and Cancer Prevention Unit of the University Hospital Vall d'Hebron. Acknowledgements to the Cellex Foundation for providing research facilities and equipment.
ICO	The authors would like to particularly acknowledge the support of the Asociación Española Contra el Cáncer (AECC), the Instituto de Salud Carlos III (organismo adscrito al Ministerio de Economía y Competitividad) and “Fondo Europeo de Desarrollo Regional (FEDER), una manera de hacer Europa” (PI10/01422, PI13/00285, PIE13/00022, PI15/00854, PI16/00563 and CIBERONC) and the Institut Català de la Salut and Autonomous Government of Catalonia (2009SGR290, 2014SGR338 and PERIS Project MedPerCan).ICO: Contract grant sponsor: Asociación Española Contra el Cáncer, Spanish Health Research Fund; Carlos III Health Institute; Catalan Health Institute and Autonomous Government of Catalonia. Contract grant numbers: ISCIII RETIC RD06/0020/1051, RD12/0036/008, PI10/01422, PI10/00748, PI13/00285, PIE13/00022, 2009SGR290 and 2014SGR364.	We wish to thank the ICO Hereditary Cancer Program team led by Dr. Gabriel Capella.

Study	Funding	Acknowledgements
IHCC	The IHCC was supported by Grant PBZ_KBN_122/P05/2004	
ILUH	The ILUH group was supported by the Icelandic Association “Walking for Breast Cancer Research” and by the Landspítali University Hospital Research Fund.	
INHERIT	This work was supported by the Canadian Institutes of Health Research for the “CIHR Team in Familial Risks of Breast Cancer” program, the Canadian Breast Cancer Research Alliance-grant #019511 and the Ministry of Economic Development, Innovation and Export Trade – grant # PSR-SIIRI-701.	We would like to thank Dr Martine Dumont, Martine Tranchant for sample management and skillful technical assistance. J.S. is Chairholder of the Canada Research Chair in Oncogenetics. J.S. and P.S. were part of the QC and Genotyping coordinating group of iCOGS (BCAC and CIMBA).
IOVCHBOCS	IOVCHBOCS is supported by Ministero della Salute and “5x1000” Istituto Oncologico Veneto grant.	
IPOBCS	This study was in part supported by Liga Portuguesa Contra o Cancro.	We wish to thank Drs. Catarina Santos, Patrícia Rocha and Pedro Pinto for their skillful contribution to the study.
KCONFAB	kConFab is supported by a grant from the National Breast Cancer Foundation, and previously by the National Health and Medical Research Council (NHMRC), the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania and South Australia, and the Cancer Foundation of Western Australia; Amanda Spurdle is supported by an NHMRC Senior Research Fellowship.	We wish to thank Heather Thorne, Eveline Niedermayr, all the kConFab research nurses and staff, the heads and staff of the Family Cancer Clinics, and the Clinical Follow Up Study (which has received funding from the NHMRC, the National Breast Cancer Foundation, Cancer Australia, and the National Institute of Health (USA)) for their contributions to this resource, and the many families who contribute to kConFab.
KOHBRA	KOHBRA is partially supported by a grant from the National R&D Program for Cancer Control, Ministry for Health, Welfare and Family Affairs, Republic of Korea (1020350 &1420190).	

Study	Funding	Acknowledgements
MAYO	MAYO is supported by NIH grants CA116167, CA128978 and CA176785, an NCI Specialized Program of Research Excellence (SPORE) in Breast Cancer (CA116201), a grant from the Breast Cancer Research Foundation, and a generous gift from the David F. and Margaret T. Grohne Family Foundation.	
MCGILL	Jewish General Hospital Weekend to End Breast Cancer, Quebec Ministry of Economic Development, Innovation and Export Trade	
MODSQUAD	MODSQUAD was supported by MH CZ - DRO (MMCI, 00209805) and by the European Regional Development Fund and the State Budget of the Czech Republic (RECAMO, CZ.1.05/2.1.00/03.0101) to LF, and by Charles University in Prague project UNCE204024 (MZ).	Modifier Study of Quantitative Effects on Disease (MODSQUAD): MODSQUAD acknowledges ModSQuaD members and Michal Zikan, Petr Pohlreich and Zdenek Kleibl (Oncogynecologic Center and Department of Biochemistry and Experimental Oncology, First Faculty of Medicine, Charles University, Prague, Czech Republic).
MUV		We wish to thank Daniela Muhr and the Senology team, and the clinicians and patients for their contributions to this study.
MSKCC	MSKCC is supported by grants from the Breast Cancer Research Foundation, the Robert and Kate Niehaus Clinical Cancer Genetics Initiative, the Andrew Sabin Research Fund. and the NIH/NCI Cancer Center Support Grant P30 CA008748.	Anne Lincoln, Lauren Jacobs
MUV		We wish to thank Daniela Muhr and the Senology team, and the clinicians and patients for their contributions to this study.
NCCS	Dr J Ngeow is supported by grants from National Medical Research Council of Singapore; Ministry of Health Health Services Research Grant Singapore and Lee Foundation Singapore	We would like to thank all patients, families and clinicians who contributed data and time to this study.
NCI	The research of Drs. MH Greene, PL Mai, and JT Loud was supported by the Intramural Research Program of the US National Cancer Institute, NIH, and by support services contracts NO2-CP-11019-50 and N02-CP-65504 with Westat, Inc, Rockville, MD.	

Study	Funding	Acknowledgements
NNPIO	This work has been supported by the Russian Federation for Basic Research (grants 14-04-93959 and 15-04-01744).	
Northshore		We would like to thank Wendy Rubinstein and the following genetic counselors for help with participant recruitment: Scott Weissman, Anna Newlin, Kristen Vogel, Lisa Dellafave-Castillo, Shelly Weiss.
NRG Oncology	This study was supported by NRG Oncology Operations grant number U10 CA180868 as well as NRG SDMC grant U10 CA180822, Gynecologic Oncology Group (GOG) Administrative Office and the GOG Tissue Bank (CA 27469) and the GOG Statistical and Data Center (CA 37517). Drs. Greene, Mai and Loud were supported by funding from the Intramural Research Program, NCI.	We thank the investigators of the Australia New Zealand NRG Oncology group
OCGN		We wish to thank members and participants in the Ontario Cancer Genetics Network for their contributions to the study.
OSU CCG	OSUCCG is supported by the Ohio State University Comprehensive Cancer Center.	Kevin Sweet, Caroline Craven, Julia Cooper, and Michelle O'Connor were instrumental in accrual of study participants, ascertainment of medical records and database management.
PBCS	This work was supported by the ITT (Istituto Toscano Tumori) grants 2011-2013.	
SEABASS	Ministry of Science, Technology and Innovation, Ministry of Higher Education (UM.C/HIR/MOHE/06) and Cancer Research Initiatives Foundation	We would like to thank Yip Cheng Har, Nur Aishah Mohd Taib, Phuah Sze Yee, Norhashimah Hassan and all the research nurses, research assistants and doctors involved in the MyBrCa Study for assistance in patient recruitment, data collection and sample preparation. In addition, we thank Philip Iau, Sng Jen-Hwei and Sharifah Nor Akmal for contributing samples from the Singapore Breast Cancer Study and the HUKM-

Study	Funding	Acknowledgements
		<p>HKL Study respectively. The Malaysian Breast Cancer Genetic Study is funded by research grants from the Malaysian Ministry of Science, Technology and Innovation, Ministry of Higher Education (UM.C/HIR/MOHE/06) and charitable funding from Cancer Research Initiatives Foundation.</p>
SMC	<p>This project was partially funded through a grant by the Israel cancer association and the funding for the Israeli Inherited breast cancer consortium</p>	<p>SMC team wishes to acknowledge the assistance of the Meirav Comprehensive breast cancer center team at the Sheba Medical Center for assistance in this study.</p>
SWE-BRCA	<p>SWE-BRCA collaborators are supported by the Swedish Cancer Society</p>	<p>Swedish scientists participating as SWE-BRCA collaborators are: from Lund University and University Hospital: Åke Borg, Håkan Olsson, Helena Jernström, Karin Henriksson, Katja Harbst, Maria Soller, Ulf Kristoffersson; from Gothenburg Sahlgrenska University Hospital: Margareta Nordling, Per Karlsson, Zakaria Einbeigi; from Stockholm and Karolinska University Hospital: Annika Lindblom, Brita Arver, Gisela Barbany Bustinza, Johanna Rantala; from Umeå University Hospital: Beatrice Melin, Christina Edwinsdotter Ardnor, Monica Emanuelsson; from Uppsala University: Maritta Hellström Pigg, Richard Rosenquist; from Linköping University Hospital: Marie Stenmark-Askmal, Sigrun Liedgren</p>

Study	Funding	Acknowledgements
UCHICAGO	UCHICAGO is supported by NCI Specialized Program of Research Excellence (SPORE) in Breast Cancer (CA125183), R01 CA142996, 1U01CA161032 and by the Ralph and Marion Falk Medical Research Trust, the Entertainment Industry Fund National Women's Cancer Research Alliance and the Breast Cancer research Foundation. OIO is an ACS Clinical Research Professor.	We wish to thank Cecilia Zvocec, Qun Niu, physicians, genetic counselors, research nurses and staff of the Cancer Risk Clinic for their contributions to this resource, and the many families who contribute to our program.
UCLA	Jonsson Comprehensive Cancer Center Foundation; Breast Cancer Research Foundation	We thank Joyce Seldon MSGC and Lorna Kwan, MPH for assembling the data for this study.
UCSF	UCSF Cancer Risk Program and Helen Diller Family Comprehensive Cancer Center	We would like to thank the following genetic counselors for participant recruitment: Beth Crawford, Kate Loranger, Julie Mak, Nicola Stewart, Robin Lee, and Peggy Conrad. And thanks to Ms. Salina Chan for her data management.
UKFOCR	UKFOCR was supported by a project grant from CRUK to Paul Pharoah.	We thank Simon Gayther, Carole Pye, Patricia Harrington and Eva Wozniak for their contributions towards the UKFOCR.
UPENN	National Institutes of Health (NIH) (R01-CA102776 and R01-CA083855; Breast Cancer Research Foundation; Susan G. Komen Foundation for the cure, Basser Research Center for BRCA	
UPITT/MWH	Frieda G. and Saul F. Shapira BRCA-Associated Cancer Research Program; Hackers for Hope Pittsburgh	
VFCTG	Victorian Cancer Agency, Cancer Australia, National Breast Cancer Foundation	Geoffrey Lindeman, Marion Harris, Martin Delatycki of the Victorian Familial Cancer Trials Group. We thank Sarah Sawyer and Rebecca Driessen for assembling this data and Ella Thompson for performing all DNA amplification.
WCP	Beth Y. Karlan was supported by the American Cancer Society Early Detection Professorship (SIOP-06-258-06-COUN) and the National Center for Advancing Translational Sciences (NCATS), Grant UL1TR000124	

Study	Funding	Acknowledgements

REFERENCES

- Abugattas J, Llacuachqui M, Allende YS, Velásquez AA, Velarde R, Cotrina J, Garcés M, León M, Calderón G, de la Cruz M and others. 2015. Prevalence of BRCA1 and BRCA2 mutations in unselected breast cancer patients from Peru. *Clin Genet* 88(4):371-5.
- Ah Mew N, Hamel N, Galvez M, Al-Saffar M, Foulkes WD. 2002. Haplotype analysis of a BRCA1: 185delAG mutation in a Chilean family supports its Ashkenazi origins. *Clinical Genetics* 62(2):151-6.
- Ahn SH, Son BH, Yoon KS, Noh DY, Han W, Kim SW, Lee ES, Park HL, Hong YJ, Choi JJ and others. 2007. BRCA1 and BRCA2 germline mutations in Korean breast cancer patients at high risk of carrying mutations. *Cancer Lett* 245(1-2):90-5.
- Aleamar B, Herzog J, Brinckmann Oliveira Netto C, Artigalás O, Schwartz IVD, Matzenbacher Bittar C, Ashton-Prolla P, Weitzel JN. 2016. Prevalence of Hispanic BRCA1 and BRCA2 mutations among hereditary breast and ovarian cancer patients from Brazil reveals differences among Latin American populations. *Cancer Genet* 209(9):417-422.
- Anczukow O, Ware MD, Buisson M, Zetoune AB, Stoppa-Lyonnet D, Sinilnikova OM, Mazoyer S. 2008. Does the nonsense-mediated mRNA decay mechanism prevent the synthesis of truncated BRCA1, CHK2, and p53 proteins? *Hum Mutat* 29(1):65-73.
- Antoniou AC, Sinilnikova OM, Simard J, Léoné M, Dumont M, Neuhausen SL, Struewing JP, Stoppa-Lyonnet D, Barjhoux L, Hughes DJ and others. 2007. RAD51 135G-->C modifies breast cancer risk among BRCA2 mutation carriers: results from a combined analysis of 19 studies. *Am J Hum Genet* 81(6):1186-200.
- Bergthorsson JT, Jonasdottir A, Johannesdottir G, Arason A, Egilsson V, Gayther S, Borg A, Hakanson S, Ingvarsson S, Barkardottir RB. 1998. Identification of a novel splice-site mutation of the BRCA1 gene in two breast cancer families: screening reveals low frequency in Icelandic breast cancer patients. *Human Mutation Suppl* 1:S195-7.
- Bernstein JL, Teraoka S, Southey MC, Jenkins MA, Andrulis IL, Knight JA, John EM, Lapinski R, Wolitzer AL, Whittemore AS and others. 2006. Population-based estimates of breast cancer risks associated with ATM gene variants c.7271T>G and c.1066-6T>G (IVS10-6T>G) from the Breast Cancer Family Registry. *Hum Mutat* 27(11):1122-8.
- Bu R, Siraj AK, Al-Obaisi KA, Beg S, Al Hazmi M, Ajarim D, Tulbah A, Al-Dayel F, Al-Kuraya KS. 2016. Identification of novel BRCA founder mutations in Middle Eastern breast cancer patients using capture and Sanger sequencing analysis. *Int J Cancer* 139(5):1091-7.
- Chen S, Parmigiani G. 2007. Meta-analysis of BRCA1 and BRCA2 penetrance. *J Clin Oncol* 25(11):1329-33.
- Chenevix-Trench G, Milne RL, Antoniou AC, Couch FJ, Easton DF, Goldgar DE, CIMBA. 2007. An international initiative to identify genetic modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: the Consortium of Investigators of Modifiers of BRCA1 and BRCA2 (CIMBA). *Breast Cancer Res* 9(2):104.
- Cock-Rada AM, Ossa CA, Garcia HI, Gomez LR. 2017. A multi-gene panel study in hereditary breast and ovarian cancer in Colombia. *Fam Cancer*.
- Domchek SM, Friebel TM, Singer CF, Evans DG, Lynch HT, Isaacs C, Garber JE, Neuhausen SL, Matloff E, Eeles R and others. 2010. Association of Risk-Reducing Surgery in BRCA1 or

- BRCA2 Mutation Carriers With Cancer Risk and Mortality. *Jama-Journal of the American Medical Association* 304(9):967-975.
- Eachkoti R, Hussain I, Afroze D, Aejazaziz S, Jan M, Shah ZA, Das BC, Siddiqi MA. 2007. BRCA1 and TP53 mutation spectrum of breast carcinoma in an ethnic population of Kashmir, an emerging high-risk area. *Cancer Lett* 248(2):308-20.
- Ferla R, Calo V, Cascio S, Rinaldi G, Badalamenti G, Carreca I, Surmacz E, Colucci G, Bazan V, Russo A. 2007. Founder mutations in BRCA1 and BRCA2 genes. *Ann Oncol* 18 Suppl 6:vi93-8.
- Friedman E, Bar-Sade Bruchim R, Kruglikova A, Risel S, Levy-Lahad E, Halle D, Bar-On E, Gershoni-Baruch R, Dagan E, Kepten I and others. 1998. Double heterozygotes for the Ashkenazi founder mutations in BRCA1 and BRCA2 genes. *Am J Hum Genet* 63(4):1224-7.
- Gao Q, Tomlinson G, Das S, Cummings S, Sveen L, Fackenthal J, Schumm P, Olopade OI. 2000. Prevalence of BRCA1 and BRCA2 mutations among clinic-based African American families with breast cancer. *Hum Genet* 107(2):186-91.
- Gayther SA, Harrington P, Russell P, Kharkevich G, Garkavtseva RF, Ponder BA. 1997. Frequently occurring germ-line mutations of the BRCA1 gene in ovarian cancer families from Russia. *Am J Hum Genet*. 60(5):1239-42.
- Gayther SA, Harrington P, Russell P, Kharkevich G, Garkavtseva RF, Ponder BA. 1997. Frequently occurring germ-line mutations of the BRCA1 gene in ovarian cancer families from Russia. *Am J Hum Genet*. 60(5):1239-42.
- Gayther SA, Mangion J, Russell P, Seal S, Barfoot R, Ponder BA, Stratton MR, Easton D. 1997. Variation of risks of breast and ovarian cancer associated with different germline mutations of the BRCA2 gene. *Nature Genetics* 15(1):103-5.
- Gayther SA, Warren W, Mazoyer S, Russell PA, Harrington PA, Chiano M, Seal S, Hamoudi R, van Rensburg EJ, Dunning AM. 1995. Germline mutations of the BRCA1 gene in breast and ovarian cancer families provide evidence for a genotype-phenotype correlation. *Nature Genetics* 11(4):428-33.
- Goldgar DE, Easton DF, Deffenbaugh AM, Monteiro AN, Tavtigian SV, Couch FJ. 2004. Integrated evaluation of DNA sequence variants of unknown clinical significance: application to BRCA1 and BRCA2. *Am J Hum Genet* 75(4):535-44.
- Gonzalez-Hormazabal P, Gutierrez-Enriquez S, Gaete D, Reyes JM, Peralta O, Waugh E, Gomez F, Margarit S, Bravo T, Blanco R and others. Spectrum of BRCA1/2 point mutations and genomic rearrangements in high-risk breast/ovarian cancer Chilean families. *Breast Cancer Res Treat*.
- Gorski B, Byrski T, Huzarski T, Jakubowska A, Menkiszak J, Gronwald J, Pluzanska A, Bebenek M, Fischer-Maliszewska L, Grzybowska E and others. 2000. Founder mutations in the BRCA1 gene in Polish families with breast-ovarian cancer. *Am J Hum Genet* 66(6):1963-8.
- Hamel N, Feng BJ, Foretova L, Stoppa-Lyonnet D, Narod SA, Imyanitov E, Sinilnikova O, Tihomirova L, Lubinski J, Gronwald J and others. 2011. On the origin and diffusion of BRCA1 c.5266dupC (5382insC) in European populations. *Eur J Hum Genet* 19(3):300-6.
- Hansen TV, Ejlersen B, Albrechtsen A, Bergsten E, Bjerregaard P, Hansen T, Myrhøj T, Nielsen PB, Timmermans-Wielenga V, Andersen MK and others. 2009. A common Greenlandic Inuit BRCA1 RING domain founder mutation. *Breast Cancer Res Treat* 115(1):69-76.
- Ho GH, Phang BH, Ng IS, Law HY, Soo KC, Ng EH. 2000. Novel germline BRCA1 mutations detected in women in singapore who developed breast carcinoma before the age of 36 years. *Cancer* 89(4):811-6.
- Jara L, Ampuero S, Santibanez E, Seccia L, Rodriguez J, Bustamante M, Martinez V, Catenaccio A, Lay-Son G, Blanco R and others. 2006. BRCA1 and BRCA2 mutations in a South American population. *Cancer Genet Cytogenet* 166(1):36-45.
- John EM, Miron A, Gong G, Phipps AI, Felberg A, Li FP, West DW, Whittemore AS. 2007. Prevalence of pathogenic BRCA1 mutation carriers in 5 US racial/ethnic groups. *JAMA* 298(24):2869-76.
- Kadalmani K, Deepa S, Bagavathi S, Anishetty S, Thangaraj K, Gajalakshmi P. 2007. Independent origin of 185delAG BRCA1 mutation in an Indian family. *Neoplasma* 54(1):51-6.

- Kaufman B, Laitman Y, Gronwald J, Lubinski J, Friedman E. 2009. Haplotype of the C61G BRCA1 mutation in Polish and Jewish individuals. *Genet Test Mol Biomarkers* 13(4):465-9.
- Kuchenbaecker KB, Hopper JL, Barnes DR, Phillips KA, Mooij TM, Roos-Blom MJ, Jervis S, van Leeuwen FE, Milne RL, Andrieu N and others. 2017. Risks of Breast, Ovarian, and Contralateral Breast Cancer for BRCA1 and BRCA2 Mutation Carriers. *JAMA* 317(23):2402-2416.
- Kurian AW. 2010. BRCA1 and BRCA2 mutations across race and ethnicity: distribution and clinical implications. *Curr Opin Obstet Gynecol* 22(1):72-8.
- Laitman Y, Borsthein RT, Stoppa-Lyonnet D, Dagan E, Castera L, Goislard M, Gershoni-Baruch R, Goldberg H, Kaufman B, Ben-Baruch N and others. Germline mutations in BRCA1 and BRCA2 genes in ethnically diverse high risk families in Israel. *Breast Cancer Res Treat*.
- Laitman Y, Feng BJ, Zamir IM, Weitzel JN, Duncan P, Port D, Thirthagiri E, Teo SH, Evans G, Latif A and others. 2013. Haplotype analysis of the 185delAG BRCA1 mutation in ethnically diverse populations. *Eur J Hum Genet* 21(2):212-6.
- Lang GT, Shi JX, Hu X, Zhang CH, Shan L, Song CG, Zhuang ZG, Cao AY, Ling H, Yu KD and others. 2017. The spectrum of BRCA mutations and characteristics of BRCA-associated breast cancers in China: Screening of 2,991 patients and 1,043 controls by next-generation sequencing. *Int J Cancer* 141(1):129-142.
- Lee AS, Ho GH, Oh PC, Balram C, Ooi LL, Lim DT, Wong CY, Hong GS. 2003. Founder mutation in the BRCA1 gene in Malay breast cancer patients from Singapore. *Human Mutation* 22(2):178.
- Li N, Zhang X, Cai Y, Xu X, Zhang L, Pan KF, Wu LY, Wang MR. 2006. BRCA1 germline mutations in Chinese patients with hereditary breast and ovarian cancer. *Int J Gynecol Cancer* 16 Suppl 1:172-8.
- Lord CJ, Ashworth A. 2017. PARP inhibitors: Synthetic lethality in the clinic. *Science* 355(6330):1152-1158.
- Maxwell KN, Domchek SM, Nathanson KL, Robson ME. 2016. Population Frequency of Germline BRCA1/2 Mutations. *J Clin Oncol* 34(34):4183-4185.
- Moslehi R, Russo D, Phelan C, Jack E, Antman K, Narod S. 2000. An unaffected individual from a breast/ovarian cancer family with germline mutations in both BRCA1 and BRCA2. *Clinical Genetics* 57(1):70-3.
- Nanda R, Schumm LP, Cummings S, Fackenthal JD, Sveen L, Ademuyiwa F, Cobleigh M, Esserman L, Lindor NM, Neuhausen SL and others. 2005. Genetic testing in an ethnically diverse cohort of high-risk women: a comparative analysis of BRCA1 and BRCA2 mutations in American families of European and African ancestry. *JAMA* 294(15):1925-33.
- NCCN. 2017. Updates in Version 2.2017 of the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast and Ovarian Cancer.
- Neuhausen SL, Godwin AK, Gershoni-Baruch R, Schubert E, Garber J, Stoppa-Lyonnet D, Olah E, Csokay B, Serova O, Lalloo F and others. 1998. Haplotype and phenotype analysis of nine recurrent BRCA2 mutations in 111 families: results of an international study. *American Journal of Human Genetics* 62(6):1381-8.
- Neuhausen SL, Mazoyer S, Friedman L, Stratton M, Offit K, Caligo A, Tomlinson G, Cannon-Albright L, Bishop T, Kelsell D and others. 1996. Haplotype and phenotype analysis of six recurrent BRCA1 mutations in 61 families: results of an international study. *American Journal of Human Genetics* 58(2):271-80.
- Oros KK, Ghadirian P, Maugard CM, Perret C, Paredes Y, Mes-Masson AM, Foulkes WD, Provencher D, Tonin PN. 2006a. Application of BRCA1 and BRCA2 mutation carrier prediction models in breast and/or ovarian cancer families of French Canadian descent. *Clin Genet* 70(4):320-9.
- Oros KK, Leblanc G, Arcand SL, Shen Z, Perret C, Mes-Masson AM, Foulkes WD, Ghadirian P, Provencher D, Tonin PN. 2006b. Haplotype analysis suggest common founders in carriers of the recurrent BRCA2 mutation, 3398delAAAAG, in French Canadian hereditary breast and/or ovarian cancer families. *BMC Med Genet* 7:23.

- Ossa CA, Torres D. 2016. Founder and Recurrent Mutations in BRCA1 and BRCA2 Genes in Latin American Countries: State of the Art and Literature Review. *Oncologist* 21(7):832-9.
- Ostrander EA, Udler MS. 2008. The role of the BRCA2 gene in susceptibility to prostate cancer revisited. *Cancer Epidemiol Biomarkers Prev* 17(8):1843-8.
- Pal T, Permuth-Wey J, Holtje T, Sutphen R. 2004. BRCA1 and BRCA2 mutations in a study of African American breast cancer patients. *Cancer Epidemiol Biomarkers Prev* 13(11 Pt 1):1794-9.
- Palomba G, Cossu A, Friedman E, Budroni M, Farris A, Contu A, Pisano M, Balduin P, Sini MC, Tanda F and others. 2007. Origin and distribution of the BRCA2-8765delAG mutation in breast cancer. *BMC Cancer* 7(1):132.
- Peto J, Collins N, Barfoot R, Seal S, Warren W, Rahman N, Easton DF, Evans C, Deacon J, Stratton MR. 1999. Prevalence of BRCA1 and BRCA2 gene mutations in patients with early-onset breast cancer. *J Natl Cancer Inst* 91(11):943-9.
- Pisano M, Cossu A, Persico I, Palmieri G, Angius A, Casu G, Palomba G, Sarobba MG, Rocca PC, Dedola MF and others. 2000. Identification of a founder BRCA2 mutation in Sardinia. *British Journal of Cancer* 82(3):553-9.
- Pritchard CC, Mateo J, Walsh MF, De Sarkar N, Abida W, Beltran H, Garofalo A, Gulati R, Carreira S, Eeles R and others. 2016. Inherited DNA-Repair Gene Mutations in Men with Metastatic Prostate Cancer. *N Engl J Med* 375(5):443-53.
- Ramus SJ, Friedman LS, Gayther SA, Ponder BA, Bobrow L, van der Looji M, Papp J, Olah E. 1997a. A breast/ovarian cancer patient with germline mutations in both BRCA1 and BRCA2. *Nat Genet*. 15(1):14-5.
- Ramus SJ, Kote-Jarai Z, Friedman LS, van der Looij M, Gayther SA, Csokay B, Ponder BA, Olah E. 1997b. Analysis of BRCA1 and BRCA2 mutations in Hungarian families with breast or breast-ovarian cancer. *American Journal of Human Genetics* 60(5):1242-6.
- Rashid MU, Zaidi A, Torres D, Sultan F, Benner A, Naqvi B, Shakoori AR, Seidel-Renkert A, Farooq H, Narod S and others. 2006. Prevalence of BRCA1 and BRCA2 mutations in Pakistani breast and ovarian cancer patients. *Int J Cancer* 119(12):2832-9.
- Rebbeck TR, Friebel TM, Mitra N, Wan F, Chen S, Andrulis IL, Apostolou P, Arnold N, Arun BK, Barrowdale D and others. 2016. Inheritance of deleterious mutations at both BRCA1 and BRCA2 in an international sample of 32,295 women. *Breast Cancer Res* 18(1):112.
- Rebbeck TR, Lynch HT, Neuhausen SL, Narod SA, Van't Veer L, Garber JE, Evans G, Isaacs C, Daly MB, Matloff E and others. 2002. Prophylactic oophorectomy in carriers of BRCA1 or BRCA2 mutations. *N Engl J Med* 346(21):1616-22.
- Rebbeck TR, Mitra N, Wan F, Sinilnikova OM, Healey S, McGuffog L, Mazoyer S, Chenevix-Trench G, Easton DF, Antoniou AC and others. 2015. Association of type and location of BRCA1 and BRCA2 mutations with risk of breast and ovarian cancer. *JAMA* 313(13):1347-61.
- Reeves MD, Yawitch TM, van der Merwe NC, van den Berg HJ, Dreyer G, van Rensburg EJ. 2004. BRCA1 mutations in South African breast and/or ovarian cancer families: evidence of a novel founder mutation in Afrikaner families. *Int J Cancer* 110(5):677-82.
- Roa BB, Boyd AA, Volcik K, Richards CS. 1996. Ashkenazi Jewish population frequencies for common mutations in BRCA1 and BRCA2. *Nature Genetics* 14(2):185-7.
- Rodríguez AO, Llacuachqui M, Pardo GG, Royer R, Larson G, Weitzel JN, Narod SA. 2012. BRCA1 and BRCA2 mutations among ovarian cancer patients from Colombia. *Gynecol Oncol* 124(2):236-43.
- Saslow D, Boetes C, Burke W, Harms S, Leach MO, Lehman CD, Morris E, Pisano E, Schnall M, Sener S and others. 2007. American Cancer Society guidelines for breast screening with MRI as an adjunct to mammography. *CA Cancer J Clin* 57(2):75-89.
- Seong MW, Cho S, Noh DY, Han W, Kim SW, Park CM, Park HW, Kim SY, Kim JY, Park SS. 2009. Comprehensive mutational analysis of BRCA1/BRCA2 for Korean breast cancer patients: evidence of a founder mutation. *Clin Genet* 76(2):152-60.

- Sharifah NA, Nurismah MI, Lee HC, Aisyah AN, Clarence-Ko CH, Naqiyah I, Rohaizak M, Fuad I, AR AJ, Zarina AL and others. Identification of novel large genomic rearrangements at the BRCA1 locus in Malaysian women with breast cancer. *Cancer Epidemiol* 34(4):442-7.
- Solano AR, Cardoso FC, Romano V, Perazzo F, Bas C, Recondo G, Santillan FB, Gonzalez E, Abalo E, Viniegra M and others. 2017. Spectrum of BRCA1/2 variants in 940 patients from Argentina including novel, deleterious and recurrent germline mutations: impact on healthcare and clinical practice. *Oncotarget* 8(36):60487-60495.
- Song CG, Hu Z, Yuan WT, Di GH, Shen ZZ, Huang W, Shao ZM. 2005. [Mutational analysis of BRCA1 and BRCA2 genes in early-onset breast cancer patients in Shanghai]. *Zhonghua Yi Xue Za Zhi* 85(43):3030-4.
- Song CG, Hu Z, Yuan WT, Di GH, Shen ZZ, Huang W, Shao ZM. 2006. [BRCA1 and BRCA2 gene mutations of familial breast cancer from Shanghai in China]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 23(1):27-31.
- Struwing JP, Hartge P, Wacholder S, Baker SM, Berlin M, McAdams M, Timmerman MM, Brody LC, Tucker MA. 1997. The risk of cancer associated with specific mutations of BRCA1 and BRCA2 among Ashkenazi Jews. *N Engl J Med* 336(20):1401-8.
- Thompson ER, Rowley SM, Li N, McInerney S, Devereux L, Wong-Brown MW, Trainer AH, Mitchell G, Scott RJ, James PA and others. 2016. Panel Testing for Familial Breast Cancer: Calibrating the Tension Between Research and Clinical Care. *J Clin Oncol* 34(13):1455-9.
- Thorlacius S, Olafsdottir G, Tryggvadottir L, Neuhausen S, Jonasson JG, Tavgigian SV, Tulinius H, Ogmundsdottir HM, Eyfjord JE. 1996. A single BRCA2 mutation in male and female breast cancer families from Iceland with varied cancer phenotypes. *Nat Genet* 13(1):117-9.
- Toh GT, Kang P, Lee SS, Lee DS, Lee SY, Selamat S, Mohd Taib NA, Yoon SY, Yip CH, Teo SH. 2008. BRCA1 and BRCA2 germline mutations in Malaysian women with early-onset breast cancer without a family history. *PLoS ONE* 3(4):e2024.
- Tonin PM, Mes-Masson AM, Narod SA, Ghadirian P, Provencher D. 1999. Founder BRCA1 and BRCA2 mutations in French Canadian ovarian cancer cases unselected for family history. *Clinical Genetics* 55(5):318-24.
- Tonin PN, Perret C, Lambert JA, Paradis AJ, Kantemiroff T, Benoit MH, Martin G, Foulkes WD, Ghadirian P. 2001. Founder BRCA1 and BRCA2 mutations in early-onset French Canadian breast cancer cases unselected for family history. *International Journal of Cancer* 95(3):189-93.
- Torres D, Rashid MU, Gil F, Umana A, Ramelli G, Robledo JF, Tawil M, Torregrosa L, Briceno I, Hamann U. 2007. High proportion of BRCA1/2 founder mutations in Hispanic breast/ovarian cancer families from Colombia. *Breast Cancer Res Treat* 103(2):225-32.
- Troudi W, Uhrhammer N, Sibille C, Dahan C, Mahfoudh W, Bouchlaka Souissi C, Jalabert T, Chouchane L, Bignon YJ, Ben Ayed F and others. 2007. Contribution of the BRCA1 and BRCA2 mutations to breast cancer in Tunisia. *J Hum Genet* 52(11):915-20.
- Velez C, Palamara PF, Guevara-Aguirre J, Hao L, Karafet T, Guevara-Aguirre M, Pearlman A, Oddoux C, Hammer M, Burns E and others. 2012. The impact of Converso Jews on the genomes of modern Latin Americans. *Hum Genet* 131(2):251-63.
- Villarreal-Garza C, Alvarez-Gómez RM, Pérez-Plasencia C, Herrera LA, Herzog J, Castillo D, Mohar A, Castro C, Gallardo LN, Gallardo D and others. 2015a. Significant clinical impact of recurrent BRCA1 and BRCA2 mutations in Mexico. *Cancer* 121(3):372-8.
- Villarreal-Garza C, Weitzel JN, Llacuachqui M, Sifuentes E, Magallanes-Hoyos MC, Gallardo L, Alvarez-Gómez RM, Herzog J, Castillo D, Royer R and others. 2015b. The prevalence of BRCA1 and BRCA2 mutations among young Mexican women with triple-negative breast cancer. *Breast Cancer Res Treat* 150(2):389-94.
- Vogel KJ, Atchley DP, Erlichman J, Broglio KR, Ready KJ, Valero V, Amos CI, Hortobagyi GN, Lu KH, Arun B. 2007. BRCA1 and BRCA2 genetic testing in Hispanic patients: mutation prevalence and evaluation of the BRCAPRO risk assessment model. *J Clin Oncol* 25(29):4635-41.

- Weitzel JN, Clague J, Martir-Negron A, Ogaz R, Herzog J, Ricker C, Jungbluth C, Cina C, Duncan P, Unzeitig G and others. 2013. Prevalence and type of BRCA mutations in Hispanics undergoing genetic cancer risk assessment in the southwestern United States: a report from the Clinical Cancer Genetics Community Research Network. *J Clin Oncol* 31(2):210-6.
- Weitzel JN, Lagos V, Blazer KR, Nelson R, Ricker C, Herzog J, McGuire C, Neuhausen S. 2005. Prevalence of BRCA mutations and founder effect in high-risk Hispanic families. *Cancer Epidemiol Biomarkers Prev* 14(7):1666-71.
- Weitzel JN, Lagos VI, Herzog JS, Judkins T, Hendrickson B, Ho JS, Ricker CN, Lowstuter KJ, Blazer KR, Tomlinson G and others. 2007. Evidence for Common Ancestral Origin of a Recurring BRCA1 Genomic Rearrangement Identified in High-Risk Hispanic Families. *Cancer Epidemiol Biomarkers Prev*.
- Welch PL, King MC. 2001. BRCA1 and BRCA2 and the genetics of breast and ovarian cancer. *Human Molecular Genetics* 10(7):705-13.
- Whittemore AS, Gong G, John EM, McGuire V, Li FP, Ostrow KL, Dicioccio R, Felberg A, West DW. 2004. Prevalence of BRCA1 mutation carriers among U.S. non-Hispanic Whites. *Cancer Epidemiol Biomarkers Prev* 13(12):2078-83.
- Zhang B, Fackenthal JD, Niu Q, Huo D, Svein WE, DeMarco T, Adebamowo CA, Ogundiran T, Olopade OI. 2009. Evidence for an ancient BRCA1 mutation in breast cancer patients of Yoruban ancestry. *Fam Cancer* 8(1):15-22.